



Figure 5 Spear II, which is 2.30m long. The spear is shown to the left of an incomplete pelvis of a horse, and the base has been broken off. Inset shows a detail of the tip of spear II. Scale in cm.

isotope stage 7, and are at least of stage 9. Correlations of the Schöningen sequence to other areas^{12,14} suggest that they were deposited during the fourth-last interglacial, probably at the end of stage 11. Judging from the mammalian fauna, the Reinsdorf assemblage is younger than the Lower Palaeolithic site of Boxgrove (West Sussex, UK)^{19,20}.

Before our discovery, the oldest complete 'spear' known was discovered in Eemian deposits at Lehringen (Lower Saxony, Germany) in 1948 (ref. 2). Thought to be about 125 kyr old (oxygen-isotope substage 5e), this thrusting spear was recovered from between the ribs of a straight-tusked elephant, and was made from yew (*Taxus*). The Schöningen spears are probably three full interglacial/glacial cycles older than the Lehringen lance (Fig. 2).

The discovery of spears designed for throwing means that theories of the development of hunting capacities and subsistence strategies of Middle Pleistocene hominids must be revised, as well-balanced, sophisticated hunting weapons were common from an early period of the Middle Pleistocene onwards. Accordingly, meat from hunting may have provided a larger dietary contribution than has previously been acknowledged^{21,22}. The Schöningen evidence also illustrates how little is known about the 'organic' component of early hominid material culture. □

Received 27 September; accepted 23 December 1996.

- Oakley, K. P., Andrews, P., Keeley, L. H. & Clark, J. D. A reappraisal of the Clacton spearpoint. *Proc. Prehist. Soc.* **43**, 13–30 (1977).
- Thieme, H. & Veil, S. Neue Untersuchungen zum einzeitlichen Elefanten-Jagdplatz Lehringen, Ldkr. Verden. *Die Kunde N.F.* **36**, 11–58 (1985).
- Thieme, H., Maier, R. & Urban, B. Archäologische Schwerpunktuntersuchungen im Helmstedter Braunkohlerevier (ASHB). – Zum Stand der Arbeiten 1983–1986. *Archäol. Korrespondenzbl.* **17**, 445–462 (1987).
- Thieme, H., Maier, R. & Urban, B. Neue Erkenntnisse zum urgeschichtlichen Siedlungsgeschehen. *Archäol. Deutsch.* **8**(2), 26–30 (1992).
- Urban, B., Thieme, H. & Elsner, H. Biostratigraphische, quartärgeologische und urgeschichtliche Befunde aus dem Tagebau 'Schöningen', Ldkr. Helmstedt. *Z. Deutsch. Geol. Ges.* **139**, 123–154 (1988).
- Urban, B., Elsner, H., Hölzer, A., Mania, D. & Albrecht, B. Eine eem- und frühweichselzeitliche Abfolge im Tagebau Schöningen, Landkreis Helmstedt. *Eiszeit. Gegenw.* **41**, 85–99 (1991).
- Urban, B., Lenhard, R., Mania, D. & Albrecht, B. Mittelpleistozän im Tagebau Schöningen, Ldkr. Helmstedt. *Z. Deutsch. Geol. Ges.* **142**, 351–372 (1991).
- Thieme, H. & Maier, R. *Archäologische Ausgrabungen im Braunkohlentagebau Schöningen, Landkreis Helmstedt.* (Hahnsche, Hannover, 1995).
- Thieme, H. & Mania, D. 'Schöningen 12'—ein mittelpleistozänes Interglazialvorkommen im Nordharzvorland mit paläolithischen Funden. *Ethnogr.-Archäol. Z.* **34**, 610–619 (1993).

- Urban, B. Mittelpleistozäne Interglaziale im Tagebau Schöningen. *Ethnogr.-Archäol. Z.* **34**, 620–622 (1993).
- Urban, B. in *Archäologische Ausgrabungen im Braunkohlentagebau Schöningen, Landkreis Helmstedt* (eds Thieme, H. & Maier, R.) 44–56 (Hahnsche, Hannover, 1995).
- Mania, D. Die Terrassen-Travertin-Sequenz von Bilzingsleben. Ein Beitrag zur Stratigraphie des Mittel- und Jungpleistozäns im Elbe-Saale-Gebiet. *Ethnogr.-Archäol. Z.* **34**, 554–575 (1993).
- Mania, D. in *Archäologische Ausgrabungen im Braunkohlentagebau Schöningen, Landkreis Helmstedt* (eds Thieme, H. & Maier, R.) 33–43 (Hahnsche, Hannover, 1995).
- Mania, D. in *The Earliest Occupation of Europe* (eds Roebroeks, W. & van Kolfschoten, T.) 85–101 (University of Leiden, 1995).
- Thieme, H., Mania, D., Urban, B. & van Kolfschoten, T. Schöningen (Nordharzvorland). Eine altpaläolithische Fundstelle aus dem mittleren Eiszeitalter. *Archäol. Korrespondenzbl.* **23**, 147–163 (1993).
- van Kolfschoten, T. Die Vertebraten des Interglazials von Schöningen 12. *Ethnogr.-Archäol. Z.* **34**, 623–628 (1993).
- van Kolfschoten, T. in *Archäologische Ausgrabungen im Braunkohlentagebau Schöningen, Landkreis Helmstedt* (eds Thieme, H. & Maier, R.) 85–94 (Hahnsche, Hannover, 1995).
- Schoch, W. H. in *Archäologische Ausgrabungen im Braunkohlentagebau Schöningen, Landkreis Helmstedt* (eds Thieme, H. & Maier, R.) 73–84 (Hahnsche, Hannover, 1995).
- Roberts, M. B., Stringer, C. B. & Parfitt, S. A. *Nature* **369**, 311–313 (1994).
- Roebroeks, W. & van Kolfschoten, T. in *The Earliest Occupation of Europe* (eds Roebroeks, W. & van Kolfschoten, T.) 297–315 (University of Leiden, 1995).
- Nitecki, M. H. in *The Evolution of Human Hunting* (eds Nitecki, M. H. & Nitecki, D. V.) 1–9 (Plenum, New York, 1987).
- Gamble, C. in *The Pleistocene Old World. Regional Perspectives* (ed. Soffer, O.) 81–98 (Plenum, New York, 1987).

Acknowledgements. We thank the Braunschweigische Kohlen-Bergwerke AG for the technical and organizational support of the archaeological excavations in the Schöningen mine; W. H. Schoch for identifying the wood; D. Mania for the drawings in Figs 1 and 2; P. Pfarr and C. S. Fuchs for the photographs in Figs 3–5; and M. Roberts, W. Roebroeks and T. van Kolfschoten for translating and reviewing the original German manuscript. The excavation of the spear site was supported by the Deutsche Stiftung Denkmalschutz.

Correspondence and requests for materials should be addressed to the author.

Viable offspring derived from fetal and adult mammalian cells

I. Wilmut, A. E. Schnieke*, J. McWhir, A. J. Kind* & K. H. S. Campbell

Roslin Institute (Edinburgh), Roslin, Midlothian EH25 9PS, UK
* PPL Therapeutics, Roslin, Midlothian EH25 9PP, UK

Fertilization of mammalian eggs is followed by successive cell divisions and progressive differentiation, first into the early embryo and subsequently into all of the cell types that make up the adult animal. Transfer of a single nucleus at a specific stage of development, to an enucleated unfertilized egg, provided an opportunity to investigate whether cellular differentiation to that stage involved irreversible genetic modification. The first offspring to develop from a differentiated cell were born after nuclear transfer from an embryo-derived cell line that had been induced to become quiescent¹. Using the same procedure, we now report the birth of live lambs from three new cell populations established from adult mammary gland, fetus and embryo. The fact that a lamb was derived from an adult cell confirms that differentiation of that cell did not involve the irreversible modification of genetic material required for development to term. The birth of lambs from differentiated fetal and adult cells also reinforces previous speculation^{1,2} that by inducing donor cells to become quiescent it will be possible to obtain normal development from a wide variety of differentiated cells.

It has long been known that in amphibians, nuclei transferred from adult keratinocytes established in culture support development to the juvenile, tadpole stage³. Although this involves differentiation into complex tissues and organs, no development to the adult stage was reported, leaving open the question of whether a differentiated adult nucleus can be fully reprogrammed. Previously we reported the birth of live lambs after nuclear transfer from cultured embryonic cells that had been induced into quiescence. We suggested that inducing the donor cell to exit the growth phase




Figure 1 Phase-contrast photomicrograph of donor-cell populations: **a**, Embryo-derived cells (SEC1); **b**, fetal fibroblasts (BLWF1); **c**, mammary-derived cells (OME). **d**, Microsatellite analysis of recipient ewes, nuclear donor cells and lambs using four polymorphic ovine markers²². The ewes are arranged from left to right

in the same order as the lambs. Cell populations are embryo-derived (SEC1), fetal-derived (BLW1), and mammary-derived (OME), respectively. Lambs have the same genotype as the donor cells and differ from their recipient mothers.

causes changes in chromatin structure that facilitate reprogramming of gene expression and that development would be normal if nuclei are used from a variety of differentiated donor cells in similar regimes. Here we investigate whether normal development to term is possible when donor cells derived from fetal or adult tissue are induced to exit the growth cycle and enter the G0 phase of the cell cycle before nuclear transfer.

Three new populations of cells were derived from (1) a day-9 embryo, (2) a day-26 fetus and (3) mammary gland of a 6-year-old ewe in the last trimester of pregnancy. Morphology of the embryo-derived cells (Fig. 1) is unlike both mouse embryonic stem (ES) cells and the embryo-derived cells used in our previous study. Nuclear transfer was carried out according to one of our established protocols¹ and reconstructed embryos transferred into recipient ewes. Ultrasound scanning detected 21 single fetuses on day 50–60 after oestrus (Table 1). On subsequent scanning at ~14-day intervals, fewer fetuses were observed, suggesting either mis-diagnosis or

fetal loss. In total, 62% of fetuses were lost, a significantly greater proportion than the estimate of 6% after natural mating⁴. Increased prenatal loss has been reported after embryo manipulation or culture of unreconstructed embryos⁵. At about day 110 of pregnancy, four fetuses were dead, all from embryo-derived cells, and post-mortem analysis was possible after killing the ewes. Two fetuses had abnormal liver development, but no other abnormalities were detected and there was no evidence of infection.

Eight ewes gave birth to live lambs (Table 1, Fig. 2). All three cell populations were represented. One weak lamb, derived from the fetal fibroblasts, weighed 3.1 kg and died within a few minutes of birth, although post-mortem analysis failed to find any abnormality or infection. At 12.5%, perinatal loss was not dissimilar to that occurring in a large study of commercial sheep, when 8% of lambs died within 24 h of birth⁶. In all cases the lambs displayed the morphological characteristics of the breed used to derive the nucleus donors and not that of the oocyte donor (Table 2). This

Table 1 Development of embryos reconstructed with three different cell types

Cell type	No. of fused couplets (%) ^a	No. recovered from oviduct (%)	No. cultured	No. of morula/blastocyst (%)	No. of morula or blastocysts transferred [†]	No. of pregnancies/no. of recipients (%)	No. of live lambs (%) [‡]
Mammary epithelium	277 (63.8) ^a	247 (89.2)	-	29 (11.7) ^a	29	1/13 (7.7)	1 (3.4%)
Fetal fibroblast	172 (84.7) ^b	124 (86.7)	-	34 (27.4) ^b	34	4/10 (40.0)	2 (5.9%)
			24	13 (54.2) ^b	6	1/6 (16.6)	1 (16.6%) [§]
Embryo-derived	385 (82.8) ^b	231 (85.3)	-	90 (39.0) ^b	72	14/27 (51.8)	4 (5.6%)
			92	36 (39.0) ^b	15	1/5 (20.0)	0

^a As assessed 1 h after fusion by examination on a dissecting microscope. Superscripts a or b within a column indicate a significant difference between donor cell types in the efficiency of fusion ($P < 0.001$) or the proportion of embryos that developed to morula or blastocyst ($P < 0.001$).

[†] It was not practicable to transfer all morulae/blastocysts.

[‡] As a proportion of morulae or blastocysts transferred. Not all recipients were perfectly synchronized.

[§] This lamb died within a few minutes of birth.



Figure 2 Lamb number 6LL3 derived from the mammary gland of a Finn Dorset ewe with the Scottish Blackface ewe which was the recipient.

alone indicates that the lambs could not have been born after inadvertent mating of either the oocyte donor or recipient ewes. In addition, DNA microsatellite analysis of the cell populations and the lambs at four polymorphic loci confirmed that each lamb was derived from the cell population used as nuclear donor (Fig. 1). Duration of gestation is determined by fetal genotype⁷, and in all cases gestation was longer than the breed mean (Table 2). By contrast, birth weight is influenced by both maternal and fetal genotype⁸. The birth weight of all lambs was within the range for single lambs born to Blackface ewes on our farm (up to 6.6 kg) and in most cases was within the range for the breed of the nuclear donor. There are no strict control observations for birth weight after embryo transfer between breeds, but the range in weight of lambs born to their own breed on our farms is 1.2–5.0 kg, 2–4.9 kg and 3–9 kg for the Finn Dorset, Welsh Mountain and Poll Dorset genotypes, respectively. The attainment of sexual maturity in the lambs is being monitored.

Development of embryos produced by nuclear transfer depends upon the maintenance of normal ploidy and creating the conditions for developmental regulation of gene expression. These responses are both influenced by the cell-cycle stage of donor and recipient cells and the interaction between them (reviewed in ref. 9). A comparison of development of mouse and cattle embryos produced by nuclear transfer to oocytes^{10,11} or enucleated zygotes^{12,13} suggests that a greater proportion develop if the recipient is an oocyte. This may be because factors that bring about reprogramming of gene expression in a transferred nucleus are required for early development and are taken up by the pronuclei during development of the zygote.

If the recipient cytoplasm is prepared by enucleation of an oocyte at metaphase II, it is only possible to avoid chromosomal damage and maintain normal ploidy by transfer of diploid nuclei^{14,15}, but further experiments are required to define the optimum cell-cycle stage. Our studies with cultured cells suggest that there is an advantage if cells are quiescent (ref. 1, and this work). In earlier studies, donor cells were embryonic blastomeres that had not been induced into quiescence. Comparisons of the phases of the growth cycle showed that development was greater if donor cells were in mitosis¹⁶ or in the G1 (ref. 10) phase of the cycle, rather than in S or G2 phases. Increased development using donor cells in G0, G1 or mitosis may reflect greater access for reprogramming factors present in the oocyte cytoplasm, but a direct comparison of these phases in the same cell population is required for a clearer understanding of the underlying mechanisms.

Table 2 Delivery of lambs developing from embryos derived by nuclear transfer from three different donor cells types, showing gestation length and birth weight

Cell type	Breed of lamb	Lamb identity	Duration of pregnancy (days)*	Birth weight (kg)
Mammary epithelium	Finn Dorset	6LL3	148	6.6
Fetal fibroblast	Black Welsh	6LL7	152	5.6
	Black Welsh	6LL8	149	2.8
	Black Welsh	6LL9†	156	3.1
Embryo-derived	Poll Dorset	6LL1	149	6.5
	Poll Dorset	6LL2‡	152	6.2
	Poll Dorset	6LL5	148	4.2
	Poll Dorset	6LL6‡	152	5.3

* Breed averages are 143, 147 and 145 days, respectively for the three genotypes Finn Dorset, Black Welsh Mountain and Poll Dorset.

† This lamb died within a few minutes of birth.

‡ These lambs were delivered by caesarian section. Overall the nature of the assistance provided by the veterinary surgeon was similar to that expected in a commercial flock.

Together these results indicate that nuclei from a wide range of cell types should prove to be totipotent after enhancing opportunities for reprogramming by using appropriate combinations of these cell-cycle stages. In turn, the dissemination of the genetic improvement obtained within elite selection herds will be enhanced by limited replication of animals with proven performance by nuclear transfer from cells derived from adult animals. In addition, gene targeting in livestock should now be feasible by nuclear transfer from modified cell populations and will offer new opportunities in biotechnology. The techniques described also offer an opportunity to study the possible persistence and impact of epigenetic changes, such as imprinting and telomere shortening, which are known to occur in somatic cells during development and senescence, respectively.

The lamb born after nuclear transfer from a mammary gland cell is, to our knowledge, the first mammal to develop from a cell derived from an adult tissue. The phenotype of the donor cell is unknown. The primary culture contains mainly mammary epithelial (over 90%) as well as other differentiated cell types, including myoepithelial cells and fibroblasts. We cannot exclude the possibility that there is a small proportion of relatively undifferentiated stem cells able to support regeneration of the mammary gland during pregnancy. Birth of the lamb shows that during the development of that mammary cell there was no irreversible modification of genetic information required for development to term. This is consistent with the generally accepted view that mammalian differentiation is almost all achieved by systematic, sequential changes in gene expression brought about by interactions between the nucleus and the changing cytoplasmic environment¹⁷. □

Methods

Embryo-derived cells were obtained from embryonic disc of a day-9 embryo from a Poll Dorset ewe cultured as described¹, with the following modifications. Stem-cell medium was supplemented with bovine DIA/LIF. After 8 days, the explanted disc was disaggregated by enzymatic digestion and cells replated onto fresh feeders. After a further 7 days, a single colony of large flattened cells was isolated and grown further in the absence of feeder cells. At passage 8, the modal chromosome number was 54. These cells were used as nuclear donors at passages 7–9. Fetal-derived cells were obtained from an eviscerated Black Welsh Mountain fetus recovered at autopsy on day 26 of pregnancy. The head was removed before tissues were cut into small pieces and the cells dispersed by exposure to trypsin. Culture was in BHK 21 (Glasgow MEM; Gibco Life Sciences) supplemented with L-glutamine (2 mM), sodium pyruvate (1 mM) and 10% fetal calf serum. At 90% confluency, the cells were passaged with a 1 : 2

division. At passage 4, these fibroblast-like cells (Fig. 1) had modal chromosome number of 54. Fetal cells were used as nuclear donors at passages 4–6. Cells from mammary gland were obtained from a 6-year-old Finn Dorset ewe in the last trimester of pregnancy¹⁸. At passages 3 and 6, the modal chromosome number was 54 and these cells were used as nuclear donors at passage numbers 3–6.

Nuclear transfer was done according to a previous protocol¹. Oocytes were recovered from Scottish Blackface ewes between 28 and 33 h after injection of gonadotropin-releasing hormone (GnRH), and enucleated as soon as possible. They were recovered in calcium- and magnesium-free PBS containing 1% FCS and transferred to calcium-free M2 medium¹⁹ containing 10% FCS at 37°C. Quiescent, diploid donor cells were produced by reducing the concentration of serum in the medium from 10 to 0.5% for 5 days, causing the cells to exit the growth cycle and arrest in G₀. Confirmation that cells had left the cycle was obtained by staining with antiPCNA/cyclin antibody (Immuno Concepts), revealed by a second antibody conjugated with rhodamine (Dakopatts).

Fusion of the donor cell to the enucleated oocyte and activation of the oocyte were induced by the same electrical pulses, between 34 and 36 h after GnRH injection to donor ewes. The majority of reconstructed embryos were cultured in ligated oviducts of sheep as before, but some embryos produced by transfer from embryo-derived cells or fetal fibroblasts were cultured in a chemically defined medium²⁰. Most embryos that developed to morula or blastocyst after 6 days of culture were transferred to recipients and allowed to develop to term (Table 1). One, two or three embryos were transferred to each ewe depending upon the availability of embryos. The effect of cell type upon fusion and development to morula or blastocyst was analysed using the marginal model of Breslow and Clayton²¹. No comparison was possible of development to term as it was not practicable to transfer all embryos developing to a suitable stage for transfer. When too many embryos were available, those having better morphology were selected.

Ultrasound scan was used for pregnancy diagnosis at around day 60 after oestrus and to monitor fetal development thereafter at 2-week intervals. Pregnant recipient ewes were monitored for nutritional status, body condition and signs of EAE, Q fever, border disease, louping ill and toxoplasmosis. As lambing approached, they were under constant observation and a veterinary surgeon called at the onset of parturition. Microsatellite analysis was carried out on DNA from the lambs and recipient ewes using four polymorphic ovine markers²².

Received 25 November 1996; accepted 10 January 1997.

1. Campbell, K. H. S., McWhir, J., Ritchie, W. A. & Wilmut, I. Sheep cloned by nuclear transfer from a cultured cell line. *Nature* **380**, 64–66 (1996).
2. Solter, D. Lambing by nuclear transfer. *Nature* **380**, 24–25 (1996).
3. Gurdon, J. B., Laskey, R. A. & Reeves, O. R. The developmental capacity of nuclei transplanted from keratinized skin cells of adult frogs. *J. Embryol. Exp. Morph.* **34**, 93–112 (1975).
4. Quinlivan, T. D., Martin, C. A., Taylor, W. B. & Cairney, I. M. Pre- and perinatal mortality in those ewes that conceived to one service. *J. Reprod. Fert.* **11**, 379–390 (1966).
5. Walker, S. K., Heard, T. M. & Seamark, R. F. *In vitro* culture of sheep embryos without co-culture: successes and perspectives. *Therio* **37**, 111–126 (1992).
6. Nash, M. L., Hungerford, L. L., Nash, T. G. & Zinn, G. M. Risk factors for perinatal and postnatal mortality in lambs. *Vet. Rec.* **139**, 64–67 (1996).
7. Bradford, G. E., Hart, R., Quirke, J. F. & Land, R. B. Genetic control of the duration of gestation in sheep. *J. Reprod. Fert.* **30**, 459–463 (1972).
8. Walton, A. & Hammond, J. The maternal effects on growth and conformation in Shire horse–Shetland pony crosses. *Proc. R. Soc. B* **125**, 311–335 (1938).
9. Campbell, K. H. S., Loi, P., Otaegui, P. J. & Wilmut, I. Cell cycle co-ordination in embryo cloning by nuclear transfer. *Rev. Reprod.* **1**, 40–46 (1996).
10. Cheong, H.-T., Takahashi, Y. & Kanagawa, H. Birth of mice after transplantation of early-cell-cycle-stage embryonic nuclei into enucleated oocytes. *Biol. Reprod.* **48**, 958–963 (1993).
11. Prather, R. S. *et al.* Nuclear transplantation in the bovine embryo. Assessment of donor nuclei and recipient oocyte. *Biol. Reprod.* **37**, 859–866 (1987).
12. McGrath, J. & Solter, D. Inability of mouse blastomere nuclei transferred to enucleated zygotes to support development *in vitro*. *Science* **226**, 1317–1318 (1984).
13. Robl, J. M. *et al.* Nuclear transplantation in bovine embryos. *J. Anim. Sci.* **64**, 642–647 (1987).
14. Campbell, K. H. S., Ritchie, W. A. & Wilmut, I. Nuclear-cytoplasmic interactions during the first cell cycle of nuclear transfer reconstructed bovine embryos: Implications for deoxyribonucleic acid replication and development. *Biol. Reprod.* **49**, 933–942 (1993).
15. Barnes, F. L. *et al.* Influence of recipient oocyte cell cycle stage on DNA synthesis, nuclear envelope breakdown, chromosome constitution, and development in nuclear transplant bovine embryos. *Mol. Reprod. Dev.* **36**, 33–41 (1993).
16. Kwon, O. Y. & Kono, T. Production of identical sextuplet mice by transferring metaphase nuclei from 4-cell embryos. *J. Reprod. Fert. Abst. Ser.* **17**, 30 (1996).
17. Gurdon, J. B. The control of gene expression in animal development (Oxford University Press, Oxford, 1974).
18. Finch, L. M. B. *et al.* Primary culture of ovine mammary epithelial cells. *Biochem. Soc. Trans.* **24**, 369S (1996).
19. Whitten, W. K. & Biggers, J. D. Complete development *in vitro* of the preimplantation stages of the mouse in a simple chemically defined medium. *J. Reprod. Fert.* **17**, 399–401 (1968).

20. Gardner, D. K., Lane, M., Spitzer, A. & Batt, P. A. Enhanced rates of cleavage and development for sheep zygotes cultured to the blastocyst stage *in vitro* in the absence of serum and somatic cells. Amino acids, vitamins, and culturing embryos in groups stimulate development. *Biol. Reprod.* **50**, 390–400 (1994).
21. Breslow, N. E. & Clayton, D. G. Approximate inference in generalized linear mixed models. *J. Am. Stat. Assoc.* **88**, 9–25 (1993).
22. Buchanan, F. C., Littlejohn, R. P., Galloway, S. M. & Crawford, A. L. Microsatellites and associated repetitive elements in the sheep genome. *Mammal. Gen.* **4**, 258–264 (1993).

Acknowledgements. We thank A. Colman for his involvement throughout this experiment and for guidance during the preparation of this manuscript; C. Wilde for mammary-derived cells; M. Ritchie, J. Bracken, M. Malcolm-Smith, W. A. Ritchie, P. Ferrier and K. Mycock for technical assistance; D. Waddington for statistical analysis; and H. Bowran and his colleagues for care of the animals. This research was supported in part by the Ministry of Agriculture, Fisheries and Food. The experiments were conducted under the Animals (Scientific Procedures) Act 1986 and with the approval of the Roslin Institute Animal Welfare and Experiments Committee.

Correspondence should be addressed to I.W. (e-mail Ian.Wilmut@bbsrc.ac.uk).

Evidence against a dedicated system for word learning in children

Lori Markson & Paul Bloom

Department of Psychology, University of Arizona, Tucson, Arizona 85721, USA

Children can learn aspects of the meaning of a new word on the basis of only a few incidental exposures and can retain this knowledge for a long period—a process dubbed ‘fast mapping’^{1–8}. It is often maintained that fast mapping is the result of a dedicated language mechanism, but it is possible that this same capacity might apply in domains other than language learning. Here we present two experiments in which three- and four-year-old children and adults were taught a novel name and a novel fact about an object, and were tested on their retention immediately, after a 1-week delay or after a 1-month delay. Our findings show that fast mapping is not limited to word learning, suggesting that the capacity to learn and retain new words is the result of learning and memory abilities that are not specific to language.

In two experiments (study 1 and study 2), 48 three-year-old children (mean age, 3 yr 7 months), 47 four-year-old children (mean age, 4 yr 5 months) and 48 undergraduate students first participated in a training phase that lasted for about twenty minutes. This phase involved the manipulation of ten kinds of objects, four of them familiar (for example, pennies) and six of them novel (see Methods). Subjects were asked to use some of the objects to measure other objects: for instance, they were asked to use pennies to measure the circumference of a plastic disc. Children were told it was a game, and adults were told it was a game designed to teach young children how to measure.

In the course of the training phase, subjects in both study 1 and study 2 were exposed to a new word—‘koba’—used to refer to one of the six unfamiliar kinds of objects. Subjects in study 1 were also taught a new fact about one or more objects belonging to another kind. They were told that the object or objects was given to the experimenter by her uncle. Subjects in study 2 were given information about an unfamiliar object, presented visually. They watched as a sticker was placed on one of the unfamiliar objects, and were told that was where the sticker should go (see Methods).

In each of the studies, one-third of the subjects from each age group were tested for comprehension immediately after the training phase, one-third were tested after a 1-week delay (6–8 days), and one-third after a 1-month delay (28–30 days). Subjects were presented with the original array of ten items and asked to recall which object was the koba. Subjects in study 1 were also asked to recall which object was given to the experimenter by her uncle. Subjects in study 2 were handed a small sticker and instructed to put it where it should go (see Methods).