

# Structural basis for Ca<sup>2+</sup> selectivity of a voltage-gated calcium channel

Lin Tang<sup>1,2\*</sup>, Tamer M. Gamal El-Din<sup>1\*</sup>, Jian Payandeh<sup>1†</sup>, Gilbert Q. Martinez<sup>1</sup>, Teresa M. Heard<sup>1</sup>, Todd Scheuer<sup>1</sup>, Ning Zheng<sup>1,2</sup> & William A. Catterall<sup>1</sup>

**Voltage-gated calcium (Ca<sub>V</sub>) channels catalyse rapid, highly selective influx of Ca<sup>2+</sup> into cells despite a 70-fold higher extracellular concentration of Na<sup>+</sup>. How Ca<sub>V</sub> channels solve this fundamental biophysical problem remains unclear. Here we report physiological and crystallographic analyses of a calcium selectivity filter constructed in the homotetrameric bacterial Na<sub>V</sub> channel Na<sub>V</sub>Ab. Our results reveal interactions of hydrated Ca<sup>2+</sup> with two high-affinity Ca<sup>2+</sup>-binding sites followed by a third lower-affinity site that would coordinate Ca<sup>2+</sup> as it moves inward. At the selectivity filter entry, Site 1 is formed by four carboxyl side chains, which have a critical role in determining Ca<sup>2+</sup> selectivity. Four carboxyls plus four backbone carbonyls form Site 2, which is targeted by the blocking cations Cd<sup>2+</sup> and Mn<sup>2+</sup>, with single occupancy. The lower-affinity Site 3 is formed by four backbone carbonyls alone, which mediate exit into the central cavity. This pore architecture suggests a conduction pathway involving transitions between two main states with one or two hydrated Ca<sup>2+</sup> ions bound in the selectivity filter and supports a 'knock-off' mechanism of ion permeation through a stepwise-binding process. The multi-ion selectivity filter of our Ca<sub>V</sub>Ab model establishes a structural framework for understanding the mechanisms of ion selectivity and conductance by vertebrate Ca<sub>V</sub> channels.**

Ca<sup>2+</sup> ions flow through Ca<sub>V</sub> channels at a rate of ~10<sup>6</sup> ions s<sup>-1</sup>, yet Na<sup>+</sup> conductance is >500-fold lower<sup>1</sup>. Such high-fidelity, high-throughput Ca<sub>V</sub> channel performance is important in regulating intracellular processes such as contraction, secretion, neurotransmission and gene expression in many different cell types<sup>2</sup>. Because the extracellular concentration of Na<sup>+</sup> is 70-fold higher than Ca<sup>2+</sup>, these essential biological functions require Ca<sub>V</sub> channels to be highly selective for Ca<sup>2+</sup> in preference to Na<sup>+</sup>, even though Ca<sup>2+</sup> and Na<sup>+</sup> have nearly identical diameters (~2 Å). Ion selectivity of Ca<sub>V</sub> channels is proposed to result from high-affinity binding of Ca<sup>2+</sup>, which prevents Na<sup>+</sup> permeation. Fast Ca<sup>2+</sup> flux through Ca<sub>V</sub> channels is thought to use a 'knock-off' mechanism in which electrostatic repulsion between Ca<sup>2+</sup> ions within the selectivity filter overcomes tight binding of a single Ca<sup>2+</sup> ion<sup>1,3-8</sup>. Most of these mechanisms require a multi-ion pore, yet extensive mutational analyses of ion selectivity and cation block of vertebrate Ca<sub>V</sub> channels support a single high-affinity Ca<sup>2+</sup>-binding site<sup>1,9-14</sup>.

Ca<sub>V</sub> channels contain a single ion-selective pore in the centre of four homologous domains<sup>2</sup>. The central pore is lined by the transmembrane segments (S) S5 and S6 and the intervening 'Pore (P)-loop' from each domain in a four-fold pseudosymmetrical arrangement. The four voltage-sensing modules composed of S1-S4 transmembrane helices are symmetrically arranged around the central pore. Ca<sub>V</sub> channels are members of the voltage-gated ion channel superfamily and are closely related to voltage-gated Na<sup>+</sup> (Na<sub>V</sub>) channels. Three structures of homotetrameric bacterial Na<sub>V</sub> channels open the way to elucidating the structural basis for ion selectivity and conductance of vertebrate Na<sub>V</sub> and Ca<sub>V</sub> channels<sup>15-17</sup>, which probably evolved from the bacterial NaChBac family and retained similar structures and functions (Supplementary Fig. 1)<sup>18-20</sup>. Interestingly, mutation of three amino-acid residues in the selectivity filter of NaChBac is sufficient to confer Ca<sup>2+</sup> selectivity<sup>21</sup>. We introduced analogous mutations into the bacterial Na<sub>V</sub> channel Na<sub>V</sub>Ab to create Ca<sub>V</sub>Ab and carried out electrophysiological and X-ray


crystallographic analyses to determine the relative permeability of Ca<sup>2+</sup> and define ion-binding sites in the selectivity filter. Our systematic analyses of Ca<sub>V</sub>Ab and intermediate derivatives provide structural and mechanistic insights into Ca<sup>2+</sup> binding and ion permeation and suggest a conductance mechanism involving two energetically similar ion-occupancy states with one or two hydrated Ca<sup>2+</sup> ions bound.

## Structure and function of Ca<sub>V</sub>Ab

Na<sub>V</sub>Ab channels have four identical pore motifs (<sup>175</sup>TLESWSM<sup>181</sup>) that form the ion selectivity filter<sup>15</sup>. The side chains of E177 form a high-field-strength site (Site<sub>HFS</sub>) at the outer end of the filter, whereas two additional potential Na<sup>+</sup>-coordination sites, a central site (Site<sub>CEN</sub>) and an inner site (Site<sub>IN</sub>), are formed by the backbone carbonyls of L176 and T175 (ref. 15). To create Ca<sub>V</sub>Ab, E177, S178 and M181 were substituted with Asp, resulting in a mutant with the pore motif <sup>175</sup>TLDDWSD<sup>181</sup> (underlined letters indicate mutated residues). Ca<sub>V</sub>Ab was expressed in *Trichopulsia ni* cells (High5) and analysed by whole-cell voltage clamp to determine its ion selectivity. In contrast to Na<sub>V</sub>Ab, which does not conduct extracellular Ca<sup>2+</sup> ions but carries outward Na<sup>+</sup> current (Fig. 1a, b), Ca<sub>V</sub>Ab conducts inward Ca<sup>2+</sup> current in a voltage-dependent manner (Fig. 1c, d). Complete titration curves for Ca<sup>2+</sup> in the presence of Ba<sup>2+</sup> as the balancing divalent cation (see Methods) revealed inhibition of Ba<sup>2+</sup> current by low concentrations of Ca<sup>2+</sup> followed by increases in Ca<sup>2+</sup> current at higher Ca<sup>2+</sup> concentrations (Fig. 1e). These results demonstrate the anomalous mole fraction effect characteristic of vertebrate Ca<sub>V</sub> channels. Comparable experiments with Na<sup>+</sup> as the balancing cation were not possible because of the instability of the High5 cells in solutions with low divalent cation concentrations. The reversal potential for Ca<sup>2+</sup> current under bi-ionic conditions closely follows the expectation for a highly Ca<sup>2+</sup>-selective conductance (30.6 ± 2.3 mV decade<sup>-1</sup>, Fig. 1f and Supplementary Fig. 2), and Ca<sub>V</sub>Ab selects Ca<sup>2+</sup> 382-fold over Na<sup>+</sup> under our standard

<sup>1</sup>Department of Pharmacology, University of Washington, Seattle, Washington 98195, USA. <sup>2</sup>Howard Hughes Medical Institute, University of Washington, Seattle, Washington 98195, USA. †Present address: Department of Structural Biology, Genentech Inc., South San Francisco, California 94080, USA.

\*These authors contributed equally to this work.



**Figure 1 | Structure and function of the  $\text{Ca}_V\text{Ab}$  channel.** **a, b,** Outward  $\text{Na}^+$  current conducted by  $\text{Na}_V\text{Ab}$  with 10 mM extracellular  $\text{Ca}^{2+}$  and 140 mM intracellular  $\text{Na}^+$ . Holding potential,  $-100$  mV; 20-ms, 10-mV step depolarizations. **c, d,** Voltage-dependent conductance of inward  $\text{Ca}^{2+}$  current by  $\text{Ca}_V\text{Ab}$  under the same conditions. 20-ms, 5-mV step depolarizations. **e,** Biphasic anomalous mole fraction effect of increasing  $\text{Ca}^{2+}$  as indicated, with  $\text{Ba}^{2+}$  as the balancing divalent cation: 10 mM  $\text{Ba}^{2+}$  with 0 to 0.5 mM  $\text{Ca}^{2+}$ , 9.3 mM  $\text{Ba}^{2+}$  with 0.7 mM  $\text{Ca}^{2+}$ , and 0 mM  $\text{Ba}^{2+}$  with 10 mM  $\text{Ca}^{2+}$

( $n = 4-10$ ). **f,** Reversal potential ( $E_{\text{rev}}$ ) versus  $\text{Ca}^{2+}$  concentration. **g,** Relative permeability of  $\text{Ca}_V\text{Ab}$  and its derivatives as measured from bi-ionic reversal potentials.  $P_{\text{Ca}}/P_{\text{Na}}$ , blue;  $P_{\text{Ba}}/P_{\text{Na}}$ , green ( $n = 5-22$ ). **h,** Cartoon representation of the overall structure of  $\text{Ca}_V\text{Ab}$  (yellow) superimposed with  $\text{Na}_V\text{Ab}$  (slate). **i, j,** Top (**i**) and side (**j**) views of the superimposed selectivity filters of  $\text{Ca}_V\text{Ab}$  (yellow) and  $\text{Na}_V\text{Ab}$  (slate) in stick representation. The three original  $\text{Na}_V\text{Ab}$  residues (black) and substituted  $\text{Ca}_V\text{Ab}$  residues (orange) are indicated. Errors bars in **b** and **d-g** are  $\pm$  s.e.m.

recording conditions, yielding a range of the permeability ( $P$ ) ratio  $P_{\text{Ca}}:P_{\text{Na}} > 10,000$ -fold for these constructs (Fig. 1g). Intermediate  $\text{Ca}_V\text{Ab}$  derivatives with single and double Asp substitutions had progressive increases in  $\text{Ca}^{2+}$  selectivity (Fig. 1g and Supplementary Fig. 2), as observed for  $\text{NaChBac}^{21}$ . The  $^{175}\text{TLDDWSN}^{181}$  mutant has an Asn residue in place of the final Asp, as observed in one domain of mammalian  $\text{Ca}_V$  channels (Supplementary Fig. 1), and it still favours  $\text{Ca}^{2+}$  over  $\text{Na}^+$  by more than 100-fold (Fig. 1g).


We crystallized and determined the structure of  $\text{Ca}_V\text{Ab}$  and its derivatives by molecular replacement using the  $\text{Na}_V\text{Ab}$  structure (PDB code 3RVY) as the search template (Supplementary Table 1). The overall structure of  $\text{Ca}_V\text{Ab}$  is very similar to that of  $\text{Na}_V\text{Ab}$ , with a root mean squared deviation (r.m.s.d.) of 0.4 Å (Fig. 1h). However, the electrostatic potential at the outer entry to the selectivity filter is more negative for  $\text{Ca}_V\text{Ab}$  than for  $\text{Na}_V\text{Ab}$  (Supplementary Fig. 3). The three negatively charged Asp residues introduced at the selectivity filter of  $\text{Ca}_V\text{Ab}$  create a wide, short, electronegatively lined pore (6 Å diameter, 10 Å length) with no significant alteration in backbone structure with respect to  $\text{Na}_V\text{Ab}$  (Fig. 1i, j and Supplementary Fig. 4). Thus, the  $\text{Ca}^{2+}$  selectivity of  $\text{Ca}_V\text{Ab}$  is mainly determined by the side chains of the amino acids at the selectivity filter.

### $\text{Ca}^{2+}$ -binding sites in the permeation pathway

The 3.2 Å resolution structure of the mutant  $^{175}\text{TLDDWSN}^{181}$  in the presence of 10 mM  $\text{Ca}^{2+}$  reveals electron densities in the selectivity filter consistent with three  $\text{Ca}^{2+}$  ions aligned on the central axis (Fig. 2a). In the outer vestibule leading to the selectivity filter, there are two additional less-intense on-axis peaks associated with weaker surrounding densities. To confirm the identity of the bound ions, we collected X-ray diffraction data at a wavelength of 1.75 Å and calculated the  $F^+_{\text{Ca}} - F^-_{\text{Ca}}$

anomalous difference map. Two strong peaks followed by a weaker peak on the intracellular side were found in the selectivity filter along the ion-conduction pathway, verifying three binding sites for  $\text{Ca}^{2+}$  (Fig. 2b). We name these Site 1, Site 2 and Site 3 from the extracellular to the intracellular side.

The  $\text{Ca}^{2+}$  ion at Site 1 is predominantly coordinated by the carboxyl groups of D178 (Site<sub>EX</sub> in  $\text{Na}_V\text{Ab}$ ), which define a plane at the selectivity filter entrance on the extracellular side of the bound  $\text{Ca}^{2+}$  ion (Fig. 2b). The distance between the carboxyl oxygen and  $\text{Ca}^{2+}$  is about 4.0 Å. This distance suggests that the ion binds at this site in a hydrated form because the ionic diameter of  $\text{Ca}^{2+}$  is 2.28 Å, too small to interact with the carboxylate anions directly but appropriate for interaction through bound water molecules. Further into the pore, the four acidic side chains of D177 (Site<sub>HES</sub> in  $\text{Na}_V\text{Ab}$ ) are located along the wall of the selectivity filter rather than projecting into the lumen, thereby also allowing the binding of a fully hydrated  $\text{Ca}^{2+}$  ion (Fig. 2b). Different from Site 1, this central  $\text{Ca}^{2+}$ -binding site (Site 2) is surrounded by a box of four carboxylate oxygen atoms from D177 above and four backbone carbonyl oxygen atoms from L176 below (Site<sub>CEN</sub> in  $\text{Na}_V\text{Ab}$ ), with oxygen- $\text{Ca}^{2+}$  distances of 4.5 Å and 4.2 Å, respectively (Fig. 2b). At the intracellular side of the pore, the third  $\text{Ca}^{2+}$ -binding site (Site 3) is composed of one plane of four carbonyls from T175 (Site<sub>IN</sub> in  $\text{Na}_V\text{Ab}$ ), which point inward to the lumen (Fig. 2b). Here the  $\text{Ca}^{2+}$  ion lies nearly on the same plane as T175 carbonyls. The chemical environment of Site 3 hints at a lower affinity, consistent with its role in exit of  $\text{Ca}^{2+}$  from the selectivity filter into the central cavity. Throughout the selectivity filter, the oxygen- $\text{Ca}^{2+}$  coordination distances are in the range of 4.0–5.0 Å, suggesting that the bound  $\text{Ca}^{2+}$  ion is continuously stabilized in a fully hydrated state when it passes through the pore. We observed diffuse electron density and in favourable cases



**Figure 2** |  $\text{Ca}^{2+}$ -binding sites in and near the selectivity filter of  $\text{NavAb}$ ,  $\text{CavAb}$  and their derivatives. **a**, Electron density at the selectivity filter of  $^{175}\text{TLDDWSN}^{181}$  (also see Supplementary Fig. 4). The  $2F_o - F_c$  electron density map (contoured at  $2\sigma$ ) of select residues in the selectivity filter with two diagonally opposed subunits shown in sticks, the  $\text{Ca}^{2+}$  ions along the ion pathway in green spheres and water molecules in red spheres. **b**, Densities at  $\text{Ca}^{2+}$  binding sites 1 and 2 from the anomalous difference Fourier map ( $3\sigma$ ) calculated from the diffraction data of a  $^{175}\text{TLDDWSN}^{181}$  mutant crystal soaked in the presence of 5 mM  $\text{Ca}^{2+}$  and collected at 1.75 Å wavelength. The distances between  $\text{Ca}^{2+}$  and oxygen atoms (dashed lines) are about 4.0 Å at Site 1 (blue lines), 4.4 Å at Site 2 (blue and magenta lines) and 5.0 Å (magenta line) at Site 3. For clarity, the subunit closest to the viewer is not shown. **c, d**, A comparison between  $^{175}\text{TLEDWSD}^{181}$  and  $^{175}\text{TLESWSD}^{181}$  ( $\text{NavAb}$ ) highlighting the importance of Site 1 for  $\text{Ca}^{2+}$  selectivity. **e, f**, A comparison between  $^{175}\text{TLDDWSD}^{181}$  ( $\text{CavAb}$ ) and  $^{175}\text{TLEDWSD}^{181}$  highlighting the role of Site 2 in fine tuning  $\text{Ca}^{2+}$  selectivity. All structures were determined in the presence of 15 mM  $\text{Ca}^{2+}$ .

discrete water molecules surrounding the bound  $\text{Ca}^{2+}$ , consistent with the presence of an inner shell of bound waters of hydration (Supplementary Fig. 5).

Although the anomalous difference map did not resolve clear peaks at the outer vestibule beyond the selectivity filter, we interpret the two on-axis  $2F_o - F_c$  densities above the three  $\text{Ca}^{2+}$  sites as two additional  $\text{Ca}^{2+}$  ions poised to enter the pore (Fig. 2a). This assignment is supported by the surrounding eight islets of density, which probably represent eight stabilized water molecules. Just as at Site 2 in the selectivity filter, these eight water molecules appear to serve as a square antiprism cage coordinating a hydrated  $\text{Ca}^{2+}$  ion at the centre (Fig. 2a). The second  $\text{Ca}^{2+}$  ion located at the bottom of this cage is  $\sim 4.5$  Å away from the four carboxyl oxygen atoms of D178, suggesting that part of its second hydration shell is replaced by D178 before the ion enters the selectivity filter. The selectivity filter, therefore, appears to select  $\text{Ca}^{2+}$

at its mouth by recognizing the  $\text{Ca}^{2+}$ - $\text{H}_2\text{O}$  hydration complex and conduct  $\text{Ca}^{2+}$  by fitting the  $\text{Ca}^{2+}$ - $\text{H}_2\text{O}$  hydration complex into the pore. Because  $\text{Ca}^{2+}$  is more electropositive than  $\text{Na}^+$ , it should bind more tightly in the ion selectivity filter of  $\text{CavAb}$ , providing a mechanistic basis for the block of  $\text{Na}^+$  permeation by  $\text{Ca}^{2+}$  at low  $\text{Ca}^{2+}$  concentration and preferential permeation of  $\text{Ca}^{2+}$  at higher  $\text{Ca}^{2+}$  concentration (see Discussion).

### Functional roles of key selectivity filter residues


Measurements of bi-ionic reversal potentials revealed that the relative permeability of different  $\text{CavAb}$  intermediate constructs for  $\text{Ca}^{2+}$  follows the order of  $\text{CavAb}$  ( $^{175}\text{TLDDWSD}^{181}$ )  $>$   $^{175}\text{TLDDWSN}^{181}$   $>$   $^{175}\text{TLEDWSD}^{181}$   $>$   $^{175}\text{TLEDWSD}^{181}$   $>$   $^{175}\text{TLEDWSD}^{181}$   $>$   $\text{NavAb}$  ( $^{175}\text{TLESWSD}^{181}$ )  $>$   $^{175}\text{TLDDWSN}^{181}$  (Fig. 1g and Supplementary Fig. 2). A comparison of the  $\text{Ca}^{2+}$  selectivity ratios between  $^{175}\text{TLEDWSD}^{181}$  and  $^{175}\text{TLESWSD}^{181}$  ( $\text{NavAb}$ ) shows that substitution of S178 with Asp is sufficient to convert the selectivity from  $\text{Na}^+$  to  $\text{Ca}^{2+}$  with  $>100$ -fold change in  $P_{\text{Ca}}:P_{\text{Na}}$  (Fig. 1g). Placement of the Asp carboxyl side chain at this position allows for the formation of the first hydrated  $\text{Ca}^{2+}$ -binding site in the selectivity filter (Fig. 2c and Supplementary Fig. 6). By contrast, S178 in  $\text{NavAb}$  binds  $\text{Ca}^{2+}$  directly by displacing its hydration shell, which blocks conductance of both  $\text{Na}^+$  and  $\text{Ca}^{2+}$  (Fig. 2d). Therefore, formation of Site 1 for binding hydrated  $\text{Ca}^{2+}$  is both necessary and sufficient for conferring  $\text{Ca}^{2+}$  selectivity over  $\text{Na}^+$  to  $\text{NavAb}$ .

The  $\text{Ca}^{2+}$  selectivity ratio of  $\text{CavAb}$  ( $^{175}\text{TLDDWSD}^{181}$ ) is 5.5-fold higher than  $^{175}\text{TLEDWSD}^{181}$  (Fig. 1g). This functional difference reflects a role of Site 2 in adjusting  $\text{Ca}^{2+}$  selectivity. Different from the side chains of D177 in  $\text{CavAb}$  ( $^{175}\text{TLDDWSD}^{181}$ ), which interact with the  $\text{Ca}^{2+}$  ion (Fig. 2e), the carboxyl group of E177 in  $^{175}\text{TLEDWSD}^{181}$  swings away from the selectivity filter and forms a hydrogen bond with D181 and the main-chain nitrogen atoms of S180 (Fig. 2f and Supplementary Fig. 7). Site 2 in  $^{175}\text{TLEDWSD}^{181}$ , therefore, is exclusively formed by the four carbonyl oxygen atoms of L176, which conceivably leads to a lower  $\text{Ca}^{2+}$ -binding affinity and a decreased  $\text{Ca}^{2+}$  selectivity. This comparison highlights both the importance of Site 2 in supporting high  $\text{Ca}^{2+}$  selectivity and the critical role of the backbone carbonyl groups of L176 in constructing this ion-binding site.

Distinct from D177 and D178, the N181 residue of  $^{175}\text{TLDDWSN}^{181}$  lies outside of the ion-conducting pore and is not directly involved in  $\text{Ca}^{2+}$  ion coordination. In close proximity to the carboxyl groups of D178, which form a ring that lines the perimeter of the pore entryway, the side chain of N181 embraces the perimeter of the D178 ring by donating a hydrogen bond to its side-chain carboxyls (Fig. 3a). Such a structural arrangement is also found in  $\text{CavAb}$  ( $^{175}\text{TLDDWSD}^{181}$ ) (Fig. 3b), although the more electronegative environment created by the extra negatively charged residue, D181, probably attracts  $\text{Ca}^{2+}$  more strongly and confers a 4- to 5-fold higher degree of  $\text{Ca}^{2+}$  selectivity to  $\text{CavAb}$  ( $^{175}\text{TLDDWSD}^{181}$ ) in comparison to  $^{175}\text{TLDDWSN}^{181}$  (Fig. 1g and Supplementary Fig. 3).

$^{175}\text{TLDDWSD}^{181}$ , which has the hydrophobic residue M181 packed next to the D178 ring, is the only  $\text{CavAb}$  intermediate that does not conduct  $\text{Ca}^{2+}$  (Supplementary Fig. 2). The crystal structure of this mutant reveals a blocking  $\text{Ca}^{2+}$  ion tightly bound at Site 1 in a dehydrated state with an oxygen-ion distance of 2.3 Å (Fig. 3c). Superposition analysis shows few structural differences between  $^{175}\text{TLDDWSD}^{181}$  and  $^{175}\text{TLDDWSN}^{181}$ , except for the side chain of D178, which is fixed by N181 in  $^{175}\text{TLDDWSN}^{181}$  but unconstrained in  $^{175}\text{TLDDWSD}^{181}$  (Fig. 3a, c). This comparison indicates that N181 in  $^{175}\text{TLDDWSN}^{181}$  and D181 in  $\text{CavAb}$  have critical roles in engaging D178 and allowing the reversible binding of the  $\text{Ca}^{2+}$ - $\text{H}_2\text{O}$  hydration complex for active  $\text{Ca}^{2+}$  conductance. Although the subtle difference in  $\text{Ca}^{2+}$  selectivity between  $^{175}\text{TLEDWSD}^{181}$  and  $^{175}\text{TLEDWSD}^{181}$  seems to argue against this conclusion (Fig. 1g), E177 in  $^{175}\text{TLEDWSD}^{181}$  actually has a structural role equivalent to that of N181 in  $^{175}\text{TLDDWSN}^{181}$  — by pointing away from the selectivity filter lumen, E177 forms a carboxylate-carboxylate





**Figure 3 | Ion binding and block of  $\text{Ca}_V\text{Ab}$  and its derivatives.** **a, b,** Top view of Site 1 with a hydrated  $\text{Ca}^{2+}$  ion coordinated by D178 with the help of N181 and D181 in  $^{175}\text{TLDDWSN}^{181}$  and  $^{175}\text{TLDDWSD}^{181}$  ( $\text{Ca}_V\text{Ab}$ ), respectively. **c,** Binding of a dehydrated  $\text{Ca}^{2+}$  ion at Site 1 in the nonconductive  $^{175}\text{TLDDWSM}^{181}$  mutant. **d,** Coordination of a hydrated  $\text{Ca}^{2+}$  ion at Site 1 of the  $^{175}\text{TLEDWSM}^{181}$  mutant. Despite the absence of a polar residue at amino acid 181, E177 in  $^{175}\text{TLEDWSM}^{181}$  is able to hold D178 in place to allow the binding of a hydrated  $\text{Ca}^{2+}$  ion. **e, f,** Block of  $\text{Ca}^{2+}$  conductance by the

indicated concentrations of  $\text{Cd}^{2+}$  and  $\text{Mn}^{2+}$ .  $^{175}\text{TLDDWSD}^{181}$ :  $\text{IC}_{50}(\text{Cd}^{2+})$ ,  $1.7 \pm 0.04 \mu\text{M}$ ;  $\text{IC}_{50}(\text{Mn}^{2+})$ ,  $526 \pm 22 \mu\text{M}$ .  $^{175}\text{TLDDWSN}^{181}$ :  $\text{IC}_{50}(\text{Cd}^{2+})$ ,  $5.9 \pm 0.4 \mu\text{M}$ ;  $\text{IC}_{50}(\text{Mn}^{2+})$ ,  $388 \pm 7 \mu\text{M}$ . Error bars are  $\pm$  s.e.m. **g, h,** Side view of the  $\text{Cd}^{2+}$ - and  $\text{Mn}^{2+}$ -binding sites in the selectivity filter of  $\text{Ca}_V\text{Ab}$ . The anomalous difference Fourier map densities (blue mesh, contoured at  $5\sigma$ ) of the bound blocking ions are calculated using diffraction data collected at  $1.75 \text{ \AA}$  wavelength. For clarity, the residues forming the selectivity filter from the two subunits in front of and behind the plane of the drawing were removed.

pair with D178 and holds it in a conduction-competent position (Fig. 3d and Supplementary Fig. 8).

### Block of $\text{Na}_V\text{Ab}$ and $\text{Ca}_V\text{Ab}$ channels by divalent cations

$\text{Cd}^{2+}$ ,  $\text{Mn}^{2+}$  and other inorganic cations are effective blockers of  $\text{Ca}_V$  channels<sup>1</sup>. Block of  $\text{Ca}^{2+}$  conductance of  $\text{Ca}_V\text{Ab}$  by  $\text{Cd}^{2+}$  and  $\text{Mn}^{2+}$  gives  $K_i$  values of  $1.78 \mu\text{M}$  for  $\text{Cd}^{2+}$  and  $526 \mu\text{M}$  for  $\text{Mn}^{2+}$  (Fig. 3e, f, blue).  $\text{Cd}^{2+}$  has a lower affinity and  $\text{Mn}^{2+}$  has a higher affinity for block of  $^{175}\text{TLDDWSN}^{181}$  (Fig. 3e, f, red). Crystals with bound  $\text{Cd}^{2+}$  and  $\text{Mn}^{2+}$  were obtained by soaking  $\text{Ca}_V\text{Ab}$  crystals in a cryo-solution containing these heavy metal ions, and the anomalous difference map was calculated from a data set collected at  $1.75 \text{ \AA}$  wavelength. The structures show that both  $\text{Cd}^{2+}$  and  $\text{Mn}^{2+}$  bind in the selectivity filter at the central site (Site 2), which is coordinated by the side chains of the four D177 residues and the carbonyl groups of L176 (Fig. 3g, h). Locked at this site, these blocking ions would inhibit the  $\text{Ca}^{2+}$  current by competitively binding to the high-affinity site required for  $\text{Ca}^{2+}$  permeation. Another important common feature of the two blocking complexes of  $\text{Ca}_V\text{Ab}$  is the block of permeation by binding of a single divalent cation within the selectivity filter, which supports the hypothesis that at least two divalent-cation-binding sites must be located close enough to induce repulsive interactions and allow divalent cation conductance by a knock-off mechanism. Because they are smaller than  $\text{Ca}^{2+}$ , the bound  $\text{Cd}^{2+}$  ( $d = 2.18 \text{ \AA}$ ) and  $\text{Mn}^{2+}$  ( $d = 1.94 \text{ \AA}$ ) must interact with the selectivity filter through bound waters of hydration, and electron density consistent with bound waters of hydration is observed in our structures (Supplementary Fig. 5).

### Ion binding at the $\text{Ca}^{2+}$ selectivity filter


To assess the properties of the three  $\text{Ca}^{2+}$ -binding sites in the selectivity filter of  $^{175}\text{TLDDWSN}^{181}$ , we titrated the concentration of  $\text{Ca}^{2+}$  in the cryo-solution and calculated the anomalous difference maps. At low  $\text{Ca}^{2+}$  concentration, two strong peaks of approximately equal intensity are found at Site 1 and Site 2 (Supplementary Fig. 9). As the  $\text{Ca}^{2+}$  concentration is raised, the electron density of Site 2 is substantially enhanced, but the peak intensity is reduced at Site 1 and remains low at Site 3 (Supplementary Fig. 9). These results suggest that the central site has the highest affinity, whereas Site 3 is the weakest. It is probable that this titration pattern reflects independent binding of  $\text{Ca}^{2+}$  to Sites 1, 2

and 3 located in different individual molecules of  $\text{Ca}_V\text{Ab}$  at low  $\text{Ca}^{2+}$  concentration, whereas increasing concentrations of  $\text{Ca}^{2+}$  saturate Site 2 in most or all individual  $\text{Ca}_V\text{Ab}$  molecules and reduce or eliminate binding at Sites 1 and 3 by repulsion. Importantly, the two flanking sites have lower affinity than the central site, as proposed in the ‘stepwise binding model’ of  $\text{Ca}_V$  channel permeation<sup>7</sup>. In this model, the presence of flanking sites of intermediate affinity facilitates the movement of  $\text{Ca}^{2+}$  into and out of a central high-affinity site, which can result in high ion conductance, even in the limiting case where there is no repulsion between bound ions.

Consistent with high binding affinity,  $\text{Ca}^{2+}$  binds at Site 2 with its first hydration shell waters coordinated with eight oxygen atoms from the channel (Fig. 2b and Supplementary Fig. 5). By contrast,  $\text{Ca}^{2+}$  at site 1 is mainly stabilized by one plane of four carboxyl groups from D178. The distance between the  $\text{Ca}^{2+}$  ion at Site 1 and the carboxyl group of D177 at Site 2 is about  $5.5\text{--}6 \text{ \AA}$ . As the  $\text{Ca}^{2+}$  ion moves inward, this distance will be reduced enough for D177 to form a stable coordination with the moving  $\text{Ca}^{2+}$  ion. This spatial configuration suggests that the two sites are separated by a low energy barrier. The differences of negative charge between D178 and the carbonyls of T175 and the differences in the geometry of their interactions with  $\text{Ca}^{2+}$  provide a plausible explanation for the higher  $\text{Ca}^{2+}$ -binding affinity at Site 1 than Site 3.

### Ion permeation mechanism

The three  $\text{Ca}^{2+}$ -binding sites in the selectivity filter of  $^{175}\text{TLDDWSN}^{181}$  are separated by a distance of about  $4.5 \text{ \AA}$ , which would result in substantial electrostatic repulsive interactions between bound ions. As in the case of the KcsA channel<sup>22</sup>, it is energetically unfavourable for  $\text{Ca}^{2+}$  ions to occupy adjacent sites simultaneously. This leads directly to our hypothesis of two interchangeable functional states of the selectivity filter in the crystal structure (Fig. 4a, b). In State 1,  $\text{Ca}^{2+}$  ions occupy Site 1 and Site 3. In State 2, a single  $\text{Ca}^{2+}$  ion occupies Site 2. These two states might be further coupled with one of the two  $\text{Ca}^{2+}$  ions at the outer vestibule ready to enter the pore (Fig. 4c). The transition between these two states occurs either when  $\text{Ca}^{2+}$  jumps from Site 1 or 3 to Site 2 or a third ion enters on one side of the filter, causing an ion to move into Site 2. It is probable that our crystal structures reflect a mixed population of  $\text{Ca}_V\text{Ab}$  molecules in which only Site 2 is occupied by  $\text{Ca}^{2+}$  plus  $\text{Ca}_V\text{Ab}$  molecules in which Site 1 and/or Site 3 are occupied. Because of



**Figure 4 | Catalytic cycle for Ca<sup>2+</sup> conductance by Ca<sub>V</sub>Ab.** **a**, An ionic occupancy state diagram of Ca<sub>V</sub>Ab showing two proposed low energy states and the potential transitions that connect them. Each state of the selectivity filter is represented by a three-box rectangle with Sites 1–3 going from left to right. Green circles represent Ca<sup>2+</sup> ions. Note that transitions in the inner circle potentially lead to ion repulsion, which might facilitate conduction. These transitions in the inner circle are more probable than those in the outer circle, as denoted by the bold arrows. **b**, The structural basis of the ionic occupancy states depicted in the inner circle of the state diagram shown on the left. The clockwise cycle represents a path for inward flux of Ca<sup>2+</sup> ions through the selectivity filter. **c**, Coupling of extracellular Ca<sup>2+</sup>-binding sites and the three sites within the selectivity filter in the two proposed ionic occupancy states. When two Ca<sup>2+</sup> ions bind to position 1 and 3 in the filter, the entryway Ca<sup>2+</sup> ion is placed furthest from the pore (left). When one Ca<sup>2+</sup> ion binds to position 2 within the filter, the ion outside the filter is pulled closer to the pore (right).

the high concentration of Ca<sup>2+</sup> in the extracellular solution, Ca<sup>2+</sup> will prefer to enter Site 1 and the weak binding of Ca<sup>2+</sup> to Site 3 will force loss of Ca<sup>2+</sup> into the low Ca<sup>2+</sup> concentration in the cytosol. This generates a unidirectional flux of Ca<sup>2+</sup> into the cell (Fig. 4c). The three-ion-occupied state would be manifest only when the external Ca<sup>2+</sup> concentration is increased enough that the flux reaches a limiting value<sup>3</sup>. The presence of the lower-affinity Site 3 flanking the central cavity would further accelerate the flux of ions by allowing stepwise binding with relatively low chemical potential energy barriers<sup>7</sup>. The combination of ionic repulsion between Ca<sup>2+</sup> ions bound at these sites and their stepwise change in binding affinity work together to allow rapid conduction in spite of the intrinsic high affinity for Ca<sup>2+</sup> binding.

## Discussion

The mechanism underlying the dramatic difference in selectivity for Ca<sup>2+</sup> over Na<sup>+</sup> in Ca<sub>V</sub>Ab versus Na<sub>V</sub>Ab is different from the mechanisms responsible for selectivity of K<sup>+</sup> over Na<sup>+</sup> and for Ca<sup>2+</sup> block revealed by high-resolution structural studies of the NaK channel<sup>23–25</sup>. In the NaK channel, K<sup>+</sup> conductance is favoured by the presence of four binding sites formed by backbone carbonyls, rather than two or three, and structural changes in amino acid residues outside the ion selectivity filter fine-tune the electronegativity of backbone carbonyls in the selectivity filter and thereby determine the affinity for block by Ca<sup>2+</sup> at the extracellular mouth of the pore. This difference in the underlying mechanism for control of ion selectivity reflects the fundamental difference in ion permeation in Na<sub>V</sub>Ab and Ca<sub>V</sub>Ab versus K<sup>+</sup> channels. In Na<sub>V</sub>Ab and Ca<sub>V</sub>Ab, permeant ions interact with both amino acid side chains and backbone carbonyls in the ion selectivity filter primarily through waters of hydration, whereas K<sup>+</sup> channels

select their permeant ions through direct interaction of the dehydrated ions with backbone carbonyls.

Our results reveal an unexpected structural basis for Ca<sup>2+</sup> selectivity and conductance in Ca<sub>V</sub>Ab channels, in which most or all interactions of Ca<sup>2+</sup> with the pore are made through its inner shell of waters of hydration. A set of three Ca<sup>2+</sup> binding sites cooperate in a knock-off mechanism in which the selectivity filter oscillates primarily between two states with either one hydrated Ca<sup>2+</sup> bound at the central site or two hydrated Ca<sup>2+</sup> ions bound at the distal sites. The high-affinity binding of Ca<sup>2+</sup> to Sites 1 and 2 ensures that Na<sup>+</sup> and other monovalent cations cannot permeate, while the high Ca<sup>2+</sup> concentration in the extracellular solution enables unidirectional flux by driving rapid occupancy of Site 1. The ionic repulsion between Ca<sup>2+</sup> ions bound at these sites and their stepwise change in binding affinity work together to allow rapid conduction in spite of the intrinsic high affinity for Ca<sup>2+</sup> binding. Although our resolution does not allow us to see all of the waters of hydration that are implied by our structure, we do observe electron density surrounding bound Ca<sup>2+</sup> ions at Sites 1, 2 and 3 that we believe represents the inner shell of waters of hydration (Supplementary Figure 5). This electron density is blurred, as if there is a diversity of arrangements of the bound water molecules in individual Ca<sub>V</sub>Ab molecules in our crystals because their hydrogen-bonding requirements can be accommodated in multiple ways between the bound cations and their coordinating carboxyl and carbonyl oxygens that comprise Sites 1, 2 and 3. In our most favourable structure (Supplementary Fig. 5g, h), four discrete water molecules are observed at Site 3. Altogether, we believe that these images provide direct support for the conclusion that bound Ca<sup>2+</sup> ions are surrounded by an inner shell of waters of hydration that are dynamic and can easily exchange local hydrogen-bonding partners. This is a unique ion conduction mechanism, which allows high-affinity interaction of hydrated Ca<sup>2+</sup> ions while mediating their rapid movement from the extracellular vestibule, through the three ion coordination sites of the selectivity filter, through the central cavity, and finally into the cytosol.

Biophysical modelling of Ca<sup>2+</sup> permeation in vertebrate Ca<sub>V</sub> channels has led to multiple proposed mechanisms, most of which involve two or more Ca<sup>2+</sup>-binding sites, yet only a single high-affinity site that is required for both permeation and Ca<sup>2+</sup> block was identified by mutagenesis and physiological analyses<sup>1</sup>. Our results with Ca<sub>V</sub>Ab channels resolve this apparent discrepancy by showing that multiple Ca<sup>2+</sup>-binding sites are necessary for permeation, but only Site 2 binds divalent cations with sufficient affinity for block. Ca<sup>2+</sup> is conducted as a hydrated cation (Supplementary Fig. 5), consistent with the large estimated functional diameter of vertebrate Ca<sub>V</sub> channels of 6 Å (ref. 26). Detailed structure–function studies of vertebrate Ca<sub>V</sub> channels show that mutations of the four residues equivalent to E177 have distinct effects on Ca<sup>2+</sup> conductance and block, implying that domain-specific interactions with Ca<sup>2+</sup> have evolved in vertebrate four-domain Ca<sub>V</sub> channels<sup>10,11,27–29</sup>. Vertebrate Ca<sub>V</sub> channels might share similar molecular mechanisms for Ca<sup>2+</sup> permeation and selectivity despite their pseudosymmetrical four-domain configuration.

*Note added in proof:* Crystal structures of isolated pore domains of other bacterial Na<sub>V</sub> channels reveal an open pore conformation for Na<sub>V</sub>Ms (ref. 30) and a binding site for blocking Ca<sup>2+</sup> ions in Na<sub>V</sub>Ae (ref. 31), which is formed primarily by the equivalent of Ser 178 in Na<sub>V</sub>Ab.

## METHODS SUMMARY

Ca<sub>V</sub>Ab and its derivative constructs were expressed in *Trichopulsia ni* insect cells and purified using anti-Flag resin and size-exclusion chromatography, reconstituted into DMPC:CHAPSO bicelles, and crystallized over an ammonium sulphate solution containing 0.1 M Na-citrate, pH 4.75. The anomalous data sets were collected at 1.75 Å wavelength with crystals soaked in a stabilizing solution containing various concentrations of cation ions. Electrophysiological experiments were performed in *T. ni* cells using standard protocols.

**Online Content** Any additional Methods, Extended Data display items and Source Data are available in the online version of the paper; references unique to these sections appear only in the online paper.

**Received 17 May; accepted 9 October 2013.**

**Published online 24 November 2013.**

- Sather, W. A. & McCleskey, E. W. Permeation and selectivity in calcium channels. *Annu. Rev. Physiol.* **65**, 133–159 (2003).
- Catterall, W. A. Voltage-gated calcium channels. *Cold Spring Harb. Perspect. Biol.* **3**, a003947 (2011).
- Almers, W. & McCleskey, E. W. Non-selective conductance in calcium channels of frog muscle: calcium selectivity in a single-file pore. *J. Physiol. (Lond.)* **353**, 585–608 (1984).
- Almers, W., McCleskey, E. W. & Palade, P. T. A non-selective cation conductance in frog muscle membrane blocked by micromolar external calcium ions. *J. Physiol. (Lond.)* **353**, 565–583 (1984).
- Hess, P. & Tsien, R. W. Mechanism of ion permeation through calcium channels. *Nature* **309**, 453–456 (1984).
- Armstrong, C. M. & Neyton, J. Ion permeation through calcium channels. A one-site model. *Ann. NY Acad. Sci.* **635**, 18–25 (1991).
- Dang, T. X. & McCleskey, E. W. Ion channel selectivity through stepwise changes in binding affinity. *J. Gen. Physiol.* **111**, 185–193 (1998).
- Lopin, K. V., Obejero-Paz, C. A. & Jones, S. W. Evaluation of a two-site, three-barrier model for permeation in  $\text{Ca}_v3.1$  ( $\alpha 1\text{G}$ ) T-type calcium channels:  $\text{Ca}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Mg}^{2+}$ , and  $\text{Na}^+$ . *J. Membr. Biol.* **235**, 131–143 (2010).
- Heinemann, S. H., Terlau, H., Stuhmer, W., Imoto, K. & Numa, S. Calcium channel characteristics conferred on the sodium channel by single mutations. *Nature* **356**, 441–443 (1992).
- Ellinor, P. T., Yang, J., Sather, W. A., Zhang, J. F. & Tsien, R. W.  $\text{Ca}^{2+}$  channel selectivity at a single locus for high-affinity  $\text{Ca}^{2+}$  interactions. *Neuron* **15**, 1121–1132 (1995).
- Yang, J., Ellinor, P. T., Sather, W. A., Zhang, J. F. & Tsien, R. W. Molecular determinants of  $\text{Ca}^{2+}$  selectivity and ion permeation in L-type  $\text{Ca}^{2+}$  channels. *Nature* **366**, 158–161 (1993).
- Kim, M. S., Morii, T., Sun, L. X., Imoto, K. & Mori, Y. Structural determinants of ion selectivity in brain calcium channel. *FEBS Lett.* **318**, 145–148 (1993).
- Cibulsky, S. M. & Sather, W. A. The EEEE locus is the sole high-affinity  $\text{Ca}^{2+}$  binding structure in the pore of a voltage-gated  $\text{Ca}^{2+}$  channel: block by  $\text{Ca}^{2+}$  entering from the intracellular pore entrance. *J. Gen. Physiol.* **116**, 349–362 (2000).
- Cloues, R. K., Cibulsky, S. M. & Sather, W. A. Ion interactions in the high-affinity binding locus of a voltage-gated  $\text{Ca}^{2+}$  channel. *J. Gen. Physiol.* **116**, 569–586 (2000).
- Payandeh, J., Scheuer, T., Zheng, N. & Catterall, W. A. The crystal structure of a voltage-gated sodium channel. *Nature* **475**, 353–358 (2011).
- Payandeh, J., Gamal El-Din, T. M., Scheuer, T., Zheng, N. & Catterall, W. A. Crystal structure of a voltage-gated sodium channel in two potentially inactivated states. *Nature* **486**, 135–139 (2012).
- Zhang, X. *et al.* Crystal structure of an orthologue of the NaChBac voltage-gated sodium channel. *Nature* **486**, 130–134 (2012).
- Ren, D. *et al.* A prokaryotic voltage-gated sodium channel. *Science* **294**, 2372–2375 (2001).
- Yu, F. H. & Catterall, W. A. The VGL-chanome: a protein superfamily specialized for electrical signaling and ionic homeostasis. *Sci. STKE* **2004**, re15 (2004).
- Koishi, R. *et al.* A superfamily of voltage-gated sodium channels in bacteria. *J. Biol. Chem.* **279**, 9532–9538 (2004).
- Yue, L., Navarro, B., Ren, D., Ramos, A. & Clapham, D. E. The cation selectivity filter of the bacterial sodium channel, NaChBac. *J. Gen. Physiol.* **120**, 845–853 (2002).
- Morais-Cabral, J. H., Zhou, Y. & MacKinnon, R. Energetic optimization of ion conduction rate by the  $\text{K}^+$  selectivity filter. *Nature* **414**, 37–42 (2001).
- Alam, A. & Jiang, Y. Structural analysis of ion selectivity in the NaK channel. *Nature Struct. Mol. Biol.* **16**, 35–41 (2009).
- Alam, A., Shi, N. & Jiang, Y. Structural insight into  $\text{Ca}^{2+}$  specificity in tetrameric cation channels. *Proc. Natl Acad. Sci. USA* **104**, 15334–15339 (2007).
- Derebe, M. G. *et al.* Tuning the ion selectivity of tetrameric cation channels by changing the number of ion binding sites. *Proc. Natl Acad. Sci. USA* **108**, 598–602 (2011).
- McCleskey, E. W. & Almers, W. The Ca channel in skeletal muscle is a large pore. *Proc. Natl Acad. Sci. USA* **82**, 7149–7153 (1985).
- Chen, X. H. & Tsien, R. W. Aspartate substitutions establish the concerted action of P-region glutamates in repeats I and III in forming the protonation site of L-type  $\text{Ca}^{2+}$  channels. *J. Biol. Chem.* **272**, 30002–30008 (1997).
- Cibulsky, S. M. & Sather, W. A. Control of ion conduction in L-type  $\text{Ca}^{2+}$  channels by the concerted action of S5–6 regions. *Biophys. J.* **84**, 1709–1719 (2003).
- Williamson, A. V. & Sather, W. A. Nonglutamate pore residues in ion selection and conduction in voltage-gated  $\text{Ca}^{2+}$  channels. *Biophys. J.* **77**, 2575–2589 (1999).
- McCusker, E. C. *et al.* Structure of a bacterial voltage-gated sodium channel pore reveals mechanisms of opening and closing. *Nature Commun.* **3**, 1102 (2012).
- Shaya, D. *et al.* Structure of a prokaryotic sodium channel pore reveals essential gating elements and an outer ion binding site common to eukaryotic channels. *J. Mol. Biol.* <http://dx.doi.org/10.1016/j.jmb.2013.10.010> (published online 10 October 2013).

**Supplementary Information** is available in the online version of the paper.

**Acknowledgements** We are grateful to the beamline staff at the Advanced Light Source (BL8.2.1 and BL8.2.2) for their assistance during data collection. Research reported in this publication was supported by the National Institute of Neurological Disorders and Stroke (NINDS) of the National Institutes of Health (NIH) under award number R01NS015751 (W.A.C.), the National Heart, Lung, and Blood Institute (NHLBI) of the NIH under award number R01HL112808 (W.A.C. and N.Z.) and a National Research Service Award from training grant T32GM008268 (T.M.H.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. This work was also supported by the Howard Hughes Medical Institute (N.Z.).

**Author Contributions** L.T., T.M.G.E.-D., J.P., T.S., N.Z. and W.A.C. designed the experiments. J.P. initiated the experimental work. L.T. conducted the protein purification, crystallization and diffraction experiments. L.T., J.P. and N.Z. determined and analysed the structures of the apo and cation-bound forms of  $\text{Ca}_v\text{Ab}$  and the intermediate  $\text{Ca}_v\text{Ab}$  constructs. T.M.G.E.-D. and T.S. performed physiological studies of  $\text{Ca}_v\text{Ab}$  and related constructs. G.Q.M. and T.M.H. made the constructs and performed the preliminary data collection. All authors interpreted the structures in light of the physiological data. L.T., N.Z. and W.A.C. wrote the manuscript with input from all co-authors. W.A.C. and N.Z. are co-senior authors.

**Author Information** Coordinates and structure factors have been deposited in the Protein Data Bank under accession codes: 4MS2 (TLDDWSN, 15 mM  $\text{Ca}^{2+}$ ), 4MTF (TLDDWSN, 0.5 mM  $\text{Ca}^{2+}$ ), 4MTG (TLDDWSN, 2.5 mM  $\text{Ca}^{2+}$ ), 4MTO (TLDDWSN, 5 mM  $\text{Ca}^{2+}$ ), 4MVM (TLDDWSN, 10 mM  $\text{Ca}^{2+}$ ), 4MVO (TLDDWSN, 15 mM  $\text{Ca}^{2+}$ ), 4MVQ (TLDDWSN, 15 mM  $\text{Ca}^{2+}$ ), 4MVR (TLDDWSN, 100 mM  $\text{Mn}^{2+}$ ), 4MVS (TLDDWSN, 100 mM  $\text{Cd}^{2+}$ ), 4MVZ (TLEDWSD, 15 mM  $\text{Ca}^{2+}$ ), 4MW3 (TLDDWSM, 15 mM  $\text{Ca}^{2+}$ ), 4MVU (TLEDWSM, 15 mM  $\text{Ca}^{2+}$ ), 4MW8 (NavAb, 15 mM  $\text{Ca}^{2+}$ ). Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints). The authors declare no competing financial interests. Readers are welcome to comment on the online version of the paper. Correspondence and requests for materials should be addressed to N.Z. (nzhang@uw.edu) or W.A.C. (wcatt@uw.edu).



## METHODS

**Protein expression and purification.** The pFastBac-Flag-Nav<sub>v</sub>Ab(I217C) that was used as the genetic background for Cav<sub>v</sub>Ab constructs was described previously<sup>15,16</sup>. Cav<sub>v</sub>Ab and its derivatives—<sup>175</sup>TLDDWSN<sup>181</sup>, <sup>175</sup>TLEDWSD<sup>181</sup>, <sup>175</sup>TLEDWSM<sup>181</sup> and <sup>175</sup>TLDSSWSM<sup>181</sup>—were generated via site-directed mutagenesis using Quick-Change (Stratagene). Recombinant baculovirus were produced using the Bac-to-Bac system (Invitrogen), and *T. ni* insect cells were infected for large-scale protein purification. Cells were collected 72 h after infection and re-suspended in buffer A (50 mM Tris-HCl, pH 8.0, 200 mM NaCl) supplemented with protease inhibitors and DNase. After sonication, digitonin (EMD Biosciences) was added to 1% and solubilization was carried out for 1–2 h at 4 °C. Clarified supernatant was then incubated with anti-Flag M2-agarose resin (Sigma) for 1–2 h at 4 °C with gentle mixing. Flag-resin was washed with ten column volumes of buffer B (buffer A supplemented with 0.12% digitonin) and eluted with buffer B supplemented with 0.1 mg ml<sup>-1</sup> Flag peptide. The eluent was concentrated and then passed over a Superdex 200 column (GE Healthcare) in 10 mM Tris-HCl, pH 8.0, 100 mM NaCl and 0.12% digitonin. The peak fractions were concentrated using a Vivaspin 30k centrifugal device.

**Crystallization and data collection.** Cav<sub>v</sub>Ab and its derivatives were concentrated to ~20 mg ml<sup>-1</sup> and reconstituted into DMPC:CHAPSO (Anatrace) bicelles according to standard protocols<sup>32,33</sup>. The protein-bicelle preparation and a well solution containing 1.8–2.0 M ammonium sulphate, 100 mM Na-citrate, pH 5.0, was mixed with a 1:1 ratio and set up in a hanging-drop vapour-diffusion format. The Ca<sup>2+</sup>-derivative crystals were obtained by soaking Cav<sub>v</sub>Ab and other mutant crystals in a cryo-protection solution (0.1 M Na-acetate, pH 5.0, 26% glucose and 2.0 M ammonium sulphate) containing the indicated concentrations of Ca<sup>2+</sup> for 40–60 min at 4 °C. The Cd<sup>2+</sup> and Mn<sup>2+</sup> derivatives were obtained by soaking Cav<sub>v</sub>Ab in the presence of 100 mM Cd<sup>2+</sup> or Mn<sup>2+</sup>, respectively. Crystals were then plunged into liquid nitrogen and maintained at 100 K during all data collection procedures.

All anomalous diffraction data sets were collected at 1.75 Å with the same synchrotron radiation source (Advanced Light Source, BL8.2.1). To optimize the anomalous signal, the data sets were collected by using the ‘inverse beam strategy’ with the wedge size of 5°.

**Structure determination, refinement and analyses.** X-ray diffraction data were integrated and scaled with the HKL2000 package<sup>34</sup> and further processed with the CCP4 package<sup>35</sup>. The structure of Cav<sub>v</sub>Ab and its derivatives were solved by molecular replacement by using an individual subunit of the Nav<sub>v</sub>Ab structure (PDB code 3RVY) as the search template. The data sets were processed in C2 space group and there are four molecules in one asymmetric unit. We chose the I222 space group to process the data sets for initial structural determination, but we found that the bound ions were slightly off-centre with respect to the axis of the pore. Therefore, to better interpret the coordination of Ca<sup>2+</sup>, Cd<sup>2+</sup> and Mn<sup>2+</sup>, we solved the structures in the C2 space group. Crystallography and NMR System software<sup>36</sup> was used for refinement of coordinates and B-factors. Final models were obtained after several cycles of refinement with REFMAC<sup>37</sup> and PHENIX<sup>38</sup> and manual re-building using COOT<sup>39</sup>. The geometries of the final structural models of Cav<sub>v</sub>Ab and its derivatives were verified using PROCHECK<sup>40</sup>. The divalent cations were identified by anomalous difference Fourier maps calculated using data collected at wavelengths of 1.75 Å for Ca<sup>2+</sup>, Cd<sup>2+</sup> and Mn<sup>2+</sup>. Detailed crystallographic data and refinement statistics for all the constructs are shown in Supplementary Table 1. All structural figures were prepared with PyMol<sup>41</sup>.

**Electrophysiology.** Wild-type NavAb expressed by infection of insect cells (High5) activates at very negative potentials ( $V_{1/2} \sim -98$  mV) and shows a strong, late use-dependent phase of slow inactivation. Mutation N49K shifts the activation curve ~75 mV to more positive potentials and abolishes the use-dependent inactivation<sup>42</sup>.

All Nav<sub>v</sub>Ab/Cav<sub>v</sub>Ab constructs used were made on the background of N49K mutation and showed good expression, allowing measurement of ionic currents 24–48 h after infection.

Whole-cell currents were recorded using an Axopatch 200 amplifier (Molecular Devices) with glass micropipettes (2–5 MΩ). Capacitance was subtracted and 80–90% of series resistance was compensated using internal amplifier circuitry. For reversal potential measurements, the intracellular pipette solution contained (in mM): 100 NaF, 10 NaCl, 20 HEPES-Na, 10 EGTA, pH 7.4 (adjusted with NaOH,  $[\text{Na}^+]_{\text{total}} = 146$  mM). Extracellular solution contained (in mM) 10 CaCl<sub>2</sub>, 140 NMDG-methanesulphonate, 20 HEPES, (pH 7.4, adjusted with Ca(OH)<sub>2</sub>,  $[\text{Ca}^{2+}]_{\text{total}} = 12$  mM). For Ba<sup>2+</sup> reversal potential measurements, BaCl<sub>2</sub> replaced CaCl<sub>2</sub>. Current–voltage (*I*–*V*) relationships were recorded in response to steps to voltages ranging from –100 to +70 mV in 5- or 10-mV increments from a holding potential of –100 mV. Pulses were generated and currents were recorded using Pulse software controlling an Instrutech ITC18 interface (HEKA). Data were analysed using Igor Pro 6.2 (WaveMetrics). Sample sizes were chosen to give s.e.m. values of less than 10% of peak values based on prior experimental experience.

Relative permeability values were calculated as described<sup>21</sup>. The permeability ratio was calculated as:

$$P_x/P_{\text{Na}} = \left\{ a_{\text{Na}} \left[ \exp\left(\frac{E_{\text{rev}}F}{RT}\right) \right] \left[ \exp\left(\frac{E_{\text{rev}}F}{RT}\right) + 1 \right] \right\} / 4a_x$$

In which *F*, *R*, *T* and *E*<sub>rev</sub> are Faraday constant, gas constant, absolute temperature and reversal potential, respectively. *a*<sub>x</sub> denotes the activity of the external divalent ion, *x*, (Ca<sup>2+</sup> or Ba<sup>2+</sup>) and *a*<sub>Na</sub>, the activity of intracellular sodium. The calculated activity coefficients were  $\gamma_{\text{Ca}} = 0.33$ ,  $\gamma_{\text{Ba}} = 0.30$ ,  $\gamma_{\text{Na}} = 0.74$ . All potentials were corrected for the experimentally determined liquid junction potential.

For anomalous mole fraction and blocking experiments, the divalent (Ca<sup>2+</sup>, Cd<sup>2+</sup> and Mn<sup>2+</sup>) was diluted in 10 mM BaCl<sub>2</sub>, 140 mM NMDG-methanesulphonate and 10 mM HEPES and perfused for 2–3 min before recording a *I*–*V* curve. The peak value of the *I*–*V* curve was measured and normalized to the peak value without the divalent cation.

- Faham, S. & Bowie, J. U. Bicelle crystallization: a new method for crystallizing membrane proteins yields a monomeric bacteriorhodopsin structure. *J. Mol. Biol.* **316**, 1–6 (2002).
- Faham, S. *et al.* Crystallization of bacteriorhodopsin from bicelle formulations at room temperature. *Protein Sci.* **14**, 836–840 (2005).
- Otwinowski, Z. & Minor, W. Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol.* **276**, 307–326 (1997).
- Collaborative Computational Project, Number 4. The CCP4 suite: programs for protein crystallography. *Acta Crystallogr. D* **50**, 760–763 (1994).
- Brünger, A. T. *et al.* Crystallography & NMR system: a new software suite for macromolecular structure determination. *Acta Crystallogr. D* **54**, 905–921 (1998).
- Murshudov, G. N., Vagin, A. A. & Dodson, E. J. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr. D* **53**, 240–255 (1997).
- Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr. D* **66**, 213–221 (2010).
- Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr. D* **60**, 2126–2132 (2004).
- Laskowski, R. A., Moss, D. S. & Thornton, J. M. Main-chain bond lengths and bond angles in protein structures. *J. Mol. Biol.* **231**, 1049–1067 (1993).
- DeLano, W. L. PyMOL molecular viewer (V.1. 2r3pre) (<http://www.pymol.org>) (2002).
- Gamal El-Din, T. M., Martinez, G. Q., Payandeh, J., Scheuer, T. & Catterall, W. A. A gating charge interaction required for late slow inactivation of the bacterial sodium channel NavAb. *J. Gen. Physiol.* **142**, 181–190 (2013).