

A Transgenomic Cytogenetic Sorghum (*Sorghum propinquum*) Bacterial Artificial Chromosome Fluorescence *in Situ* Hybridization Map of Maize (*Zea mays* L.) Pachytene Chromosome 9, Evidence for Regions of Genome Hyperexpansion

F. Ina E. Amarillo and Hank W. Bass¹

Department of Biological Science, Florida State University, Tallahassee, Florida 32306-4370

Manuscript received August 20, 2007

Accepted for publication September 23, 2007

ABSTRACT

A cytogenetic FISH map of maize pachytene-stage chromosome 9 was produced with 32 maize marker-selected sorghum BACs as probes. The genetically mapped markers used are distributed along the linkage maps at an average spacing of 5 cM. Each locus was mapped by means of multicolor direct FISH with a fluorescently labeled probe mix containing a whole-chromosome paint, a single sorghum BAC clone, and the centromeric sequence, CentC. A maize-chromosome-addition line of oat was used for bright unambiguous identification of the maize 9 fiber within pachytene chromosome spreads. The locations of the sorghum BAC-FISH signals were determined, and each new cytogenetic locus was assigned a centiMcClintock position on the short (9S) or long (9L) arm. Nearly all of the markers appeared in the same order on linkage and cytogenetic maps but at different relative positions on the two. The CentC FISH signal was localized between *cdo17* (at 9L.03) and *tda66* (at 9S.03). Several regions of genome hyperexpansion on maize chromosome 9 were found by comparative analysis of relative marker spacing in maize and sorghum. This transgenomic cytogenetic FISH map creates anchors between various maps of maize and sorghum and creates additional tools and information for understanding the structure and evolution of the maize genome.

THE genome of maize (*Zea mays* L.) has been studied as a model for eukaryotic genetics, cereal crops, and monocot genome evolution (CHANDLER and BRENDEL 2002), but its size and organizational complexity complicate resolution of its structure. The presence of large gene-poor areas, segmental duplications, abundant retrotransposons, and microvariation among lines of maize all confound efforts to develop a fully assembled physical map of the entire maize genome (KUMAR and BENNETZEN 1999; GAUT *et al.* 2000; MEYERS *et al.* 2001; YUAN *et al.* 2003; MESSING *et al.* 2004; SWIGONOVA *et al.* 2004; PATERSON *et al.* 2005). Despite these complexities, several different kinds of maps have been developed to characterize its structure and function at the DNA and chromosome levels. Many linkage maps have been developed, including those based on mutant phenotypes (EMERSON *et al.* 1935) and more recently those that include thousands of additional molecular markers such as restriction fragment length polymorphisms (RFLPs), simple sequence repeats (SSRs), single-nucleotide polymorphisms (SNPs), and insertion-deletion polymorphisms (indels) (HELENTJARIS *et al.* 1986; COE *et al.* 1987; BURR *et al.* 1988; CAUSSE *et al.*

1996; SENIOR and HEUN 1993; TARMINO and TINGEY 1996; HARUSHIMA *et al.* 1998; DAVIS *et al.* 1999; LEE *et al.* 2002; SHAROPOVA *et al.* 2002; BOWERS *et al.* 2003; FU *et al.* 2006). Physical maps of overlapping clones have been produced and anchored to the linkage map by means of molecular probes (DE JONG *et al.* 1999; BENNETZEN *et al.* 2001; CHANDLER and BRENDEL 2002; GARDINER *et al.* 2004; MESSING *et al.* 2004; BOWERS *et al.* 2005; SONG *et al.* 2005; HASS-JACOBUS *et al.* 2006). Another type of physical map is the cytological map produced by direct microscopic inspection of the chromosomes that make up the nuclear genome. Cytogenetic maps are valuable because they can place genetic loci directly within the entire chromosome, the ultimate contig, providing information on the location, order, and distribution of DNA sequences in relation to other genetic markers along the chromosomes (SADDER *et al.* 2000; ANDERSON *et al.* 2004; KIM *et al.* 2005a,b). In contrast to those of the well-developed linkage maps and clone- or sequence-based physical maps, the construction of high-density cytogenetic maps is nascent and relatively underdeveloped.

Cytological analysis of maize meiotic chromosomes provided fundamental insights into transmission genetics and the dynamic nature of the maize genome. The early insights included the physical basis of genetic recombination, the discovery of transposable DNA

¹Corresponding author: Department of Biological Science, Florida State University, Tallahassee, FL 32306-4370. E-mail: bass@bio.fsu.edu

elements, the capping properties of telomeres, and evidence in support of the chromosome theory of inheritance (CREIGHTON and McCLINTOCK 1931; RHOADES and McCLINTOCK 1935; McCLINTOCK 1941, 1978; RHOADES 1950; and reviewed by CARLSON 1988). These earlier studies also provided the basis for the meiotic chromosome karyotype of maize, in which chromosome-spread preparations allow the 10 individual maize chromosomes to be recognized. Cytological chromosome stains reveal the presence of chromosomal landmarks, such as the knobs and centromeres, but the meiotic as well as the somatic karyotype of maize has lacked extensive genetic detail for many decades (CARLSON 1988; DEMPSEY 1994; CHEN *et al.* 2000; ADAWY *et al.* 2004).

Advances in molecular biology and genomics offered new tools for cytological localization of DNA sequences and prospects for further cytogenetic map development in maize (DE JONG *et al.* 1999; HARPER and CANDE 2000; SADDER *et al.* 2000). The development of cytogenetic FISH maps of maize has progressed from mapping repeat sequences such as knobs, centromeres, and telomeres to mapping RFLP markers, single-copy genes, and individual transposons on mitotic and meiotic chromosomes (SHEN *et al.* 1987; COE 1994; CHEN *et al.* 2000; SADDER *et al.* 2000; SADDER and WEBER 2001, 2002; KOUMBARIS and BASS 2003; KATO *et al.* 2004, 2005; WANG *et al.* 2006; LAMB *et al.* 2007). FISH mapping of pachytene chromosomes has proved to be useful for many other plant species such as tomato (ZHONG *et al.* 1996a,b; PETERSON *et al.* 1999), potato (SONG *et al.* 2000), *Arabidopsis* (FRANSZ *et al.* 1996; LYSAK *et al.* 2001), *Medicago* (KULIKOVA *et al.* 2001), rice (CHENG *et al.* 2001a,b, 2002), sorghum (ISLAM-FARIDI *et al.* 2002; KIM *et al.* 2005a,b), Brassica (HOWELL *et al.* 2002), and soybean (WALLING *et al.* 2006).

Mitotic and meiotic chromosomes have been successfully used to create cytogenetic maps of maize. The mitotic chromosomes are easier to prepare, but meiotic chromosomes have the advantage of longer axial fibers for improved localization within chromosome arms (PEDERSEN and LINDE-LAURSEN 1994; CHENG *et al.* 2001a,b; DESEL *et al.* 2001; KULIKOVA *et al.* 2001). Another advantage of meiotic chromosomes is that the pachytene-based cytogenetic maps can be compared to and integrated directly with recombination nodule-based maps and translocation breakpoint data (ANDERSON *et al.* 2004; SHERIDAN and AUGER 2006).

Major challenges for any FISH-based mapping technique include the detection of small gene-size fragments, target chromosome identification, and probe specificity. Probe-detection limits can be overcome if large insert clones are used, such as those carried in BAC, YAC, or cosmid vectors (WOO *et al.* 1994; HANSON *et al.* 1995; JIANG *et al.* 1995; OHMIDO *et al.* 1998; ZWICK *et al.* 1998; ZHONG *et al.* 1999; DONG *et al.* 2000; KULIKOVA *et al.* 2001). The advantage of increased signal strength with increasing probe size is offset, however, by the

commensurate increase in the likelihood of detecting unintended targets such as repetitive sequences. This problem is acute in maize, where intergenic repetitive sequence elements abound and any given maize BAC clone may only contain a few kilobase pairs of unique, single-copy sequence (LIU *et al.* 2007). One approach to FISH mapping in maize is to use large maize genomic DNA fragments in conjunction with competitive *in situ* suppression hybridization (SADDER *et al.* 2000; SADDER and WEBER 2002). Another approach is to find relatively large single-gene fragments (>3 kbp) for loci to be mapped (WANG *et al.* 2006). Yet another strategy is to use genomic BAC clones from the small-genome relative sorghum (KOUMBARIS and BASS 2003). The cross-hybridization of DNA probes from one species to target chromosomes of another species can help overcome detection limits if the two species have different or divergent classes of interspersed repetitive sequences, as is the case for sorghum and maize. Transgenomic mapping and large-fragment FISH have been successfully used in maize and in other plant species for comparative genomics (HULBERT *et al.* 1990; FUCHS *et al.* 1996; GOMEZ *et al.* 1997; ZWICK *et al.* 1998; JACKSON *et al.* 2000; BOWERS *et al.* 2003; KOUMBARIS and BASS 2003).

KOUMBARIS and BASS (2003) developed a technique combining transgenomic and BAC-FISH mapping to overcome the probe-detection limit and establish an indirect way to define the cytogenetic location of sequences corresponding to targets such as RFLP probes, many of which are <1 kb. The method takes advantage of genomic and genetic resources in maize and sorghum. Here we report results from extension of this technique to production of the first detailed transgenomic BAC-FISH map of any maize chromosome. The locations of 32 genetically mapped markers were placed on the cytogenetic map, and their distribution revealed distinct irregularities with implications for maize genome assembly and evolution.

MATERIALS AND METHODS

Meiotic chromosome spreads: Chromosome spreads were prepared from a disomic maize chromosome-addition line of oat (OMAd9.2b from KYNAST *et al.* 2001), referred to as "oat-maize 9" in this study, that was grown in plant growth chambers or in a greenhouse (Mission Road Facility, Biological Science, Florida State University, Tallahassee, FL). Meiosis-stage florets were harvested and fixed in Carnoy's solution (3 parts absolute ethanol:1 part glacial acetic acid) for 1–2 days on a rotatory shaker at 4°. After fixation, the florets were rinsed with distilled water and stored in 70% ethanol at –20° until used. The meiotic stage was determined for one of the three anthers from a floret by the acetocarmine method. Anthers from florets with pachytene-stage meiocytes were collected and stored in 70% ethanol at –20° until used. Anthers were digested with enzymes as described by ZHONG *et al.* (1996b), and during pachytene spread preparation, slides were given an extra three to five rounds of water vapor-acetic

acid exposure, which allowed more spreading of pachytene chromosomes (Koumbaris and Bass 2003). The quality of spreads was evaluated by differential interference contrast microscopy. Slides with numerous well-spread chromosomes located in the middle of the slide, with good chromosome morphology and minimal cellular debris, were stored at -20° until used for FISH.

Identification and selection of sorghum BAC clones:

Sorghum BAC clones used as FISH probes were screened by hybridization with maize RFLP probes (as described below and by Koumbaris and Bass 2003) or with overgo probes (carried out by Bowers *et al.* 2005) designed to detect various maize marker probes. For RFLP-selected BACs, *Sorghum propinquum* genomic BAC grid-library filter arrays were obtained from A. H. Paterson (YRL filter pair; University of Georgia, Athens, GA). The detected *S. propinquum* BAC clones (typically four to seven overlapping BACs for each RFLP probe) were grown and the BAC DNA was initially isolated by the plasmid miniprep method with the QIAprep Spin Miniprep kit (no. 27104; QIAGEN, Valencia, CA). Southern blot analysis was then used on the restriction enzyme-digested miniprep DNA to verify clone identity. The criteria for BAC clone selection were (1) the BAC should belong to a group of overlapping clones detected by the probe and within one contig, (2) the BAC should be centrally located within this group of probe-detected overlapping BACs, and (3) the BAC should contain the same size restriction enzyme fragment as that observed with most or all of the other probe-detected overlapping BAC clones. A single selected sorghum BAC clone for each locus was then grown for large-scale BAC DNA preparation with the QIAGEN Large Construct kit (no. 12462), according to manufacturer's instructions. Highly purified BAC DNA was digested with *EcoRI* enzyme and then direct labeled for FISH by random-primed labeling. The YRL filters were reused after stripping by two to three washes for 30 min each with 15 mM sodium phosphate buffer (pH 6.9) at 80° . In addition to these BACs, we obtained others as gifts, including a *S. bicolor* BAC corresponding to the maize waxy1 locus (BAC 131L1, GenBank accession AF488412), from J. Ma and J. Bennetzen and several other *S. bicolor* BACs (sbb18256/191b4 and sb16685/174g5) that had been used as FISH probes on pachytene chromosomes (Islam-Faridi *et al.* 2002) from P. E. Klein (Texas A&M University, College Station, TX).

Slide pretreatment and FISH with pachytene chromosome spreads: The overall FISH procedure was performed as described by Koumbaris and Bass (2003) with minor modifications. Pachytene chromosomes were denatured in 70% formamide in $2\times$ SSC at 70° for 3–5 min and then dehydrated for 3 min in each concentration of an ice-cold ethanol series (70, 90, and 100%). The BAC clones were labeled with ChromaTide Alexa Fluor 546-16-OBEA-dCTP (Invitrogen, San Diego). The labeled FISH probes were concentrated by ethanol precipitation, redissolved in TE, and stored at -20° until used. The probe mix was denatured for 10 min at 90° , quick cooled on ice, and then combined with formamide, SSC, and dextran sulfate. The final FISH probe mixture consisted of 100 μ g/ml Alexa fluor 488-labeled maize DNA (inbred KWF), centromere-specific CentC probe (10 μ g/ml Alexa 647-labeled CentC repeat (Ananiev *et al.* 1998) DNA or 5–10 μ g/ml of the oligonucleotide probe MCCY (labeled with Cy5), 200–300 μ g/ml Alexa 546-labeled sorghum BAC DNA, 1 μ g/ml calf thymus DNA, and 180 μ g/ml *S. bicolor* (genotype Tx623) genomic DNA in $2\times$ SSC, 50% formamide, and 10% dextran sulfate. The probe mix was added to the slide bearing the target pachytene chromosomes. The SSC content was decreased to as little as $0.5\times$ SSC in cases where high background was initially observed. Hybridization was carried out at 37° on slides (20–25 μ l per slide) with rubber cement-sealed cover-

slips with the twin tower block of the DNA engine tetrad (PTC-225; MJ Research, Watertown, MA) for 18–20 hr. After hybridization, slides were washed three times (5 min each) with 50% formamide plus $0.5-2\times$ SSC at 37° and then washed at room temperature with $2\times$ SSC three times (5 min each), washed with $1\times$ PBS three times (5 min each), stained with 3–5 μ g/ml DAPI, washed with 1 mM DTT in $1\times$ PBS, and then mounted with Vectashield (Vector Labs, Burlingame, CA) for microscopy.

Data collection and image processing: The FISH preparations were analyzed with an Olympus microscope equipped with a CCD camera (Applied Precision, Issaquah, WA). Three-dimensional (3D) images of oat-maize chromosome fibers, maize pachytene chromosome 9, sorghum BAC-FISH signal, and maize centC MCCY signal were acquired on the DAPI, FITC, RHOD, and Cy5 channels, respectively. The 3D image stacks (with Z sections spaced at 0.2–0.3 μ m for a total depth of 5–8 μ m) were subjected to iterative 3D deconvolution, and nuclei showing well-stained maize pachytene fibers were used to trace and computationally straighten maize 9 chromosomes for cytogenetic mapping. Pachytene fibers with little or no background staining were straightened and analyzed as described in Koumbaris and Bass (2003).

FISH locus determination and nomenclature: The arm of interest was divided into 20–40 bins of equal length, and any FISH signal on the arm was assigned to a bin on the basis of its fractional distance from the centromere (position 0.0) to the telomere (position 1.0). Frequency histograms were inspected to identify regions with significant, above-background signals as described by Koumbaris and Bass (2003). The measured positions for all FISH signals that fell within these peak regions were averaged. The mean value, standard error (SEM), and sample size (n) are given in Table 1 for each locus in centiMcClintock (cMC) units. The nomenclature system (as described in <http://www.maizegdb.org/CMMprotocols.php>) for a FISH locus consists of (1) the sorghum BAC source (sbb for *S. bicolor* or spb for *S. propinquum*), (2) location on the linkage map (CBM for core bin marker or just the resident bin number for other markers), and (3) the cytogenetic map position, followed by (4) the maize RFLP marker name in parentheses.

RESULTS

Selection of maize markers for cytogenetic mapping:

A detailed pachytene-stage cytogenetic FISH map of maize chromosome 9 was created with sorghum BACs that correspond to well-characterized maize RFLP probes. Maize RFLP probes have been used in hundreds of linkage studies but most of these probes are smaller than the lower limit of FISH detection on meiotic chromosomes as recently determined by Wang *et al.* (2006). We used an indirect mapping strategy that overcomes this detection limit by treating maize marker-selected syntenic sorghum BACs as surrogate FISH probes. This strategy was initially described by Koumbaris and Bass (2003) for three loci and is extended here for an additional 32 cytogenetic loci.

We initially selected 52 markers that are spaced ~ 5 cM apart covering the entire linkage map, as shown in Figure 1, using the University of Missouri-Columbia (UMC) 98 linkage map as our base linkage map (Davis *et al.* 1999). This high-density map is saturated with maize




FIGURE 1.—Location of RFLP markers chosen for cytogenetic FISH mapping and their position on the linkage map of maize chromosome 9. (A) Partial linkage map of chromosome 9 adapted from UMC 98 9 (DAVIS *et al.* 1999). Core bin markers (CBM) are in boxes and their corresponding cumulative map position in centimorgans is indicated on the left. The genetic bins (9.00, 9.01, etc.) are indicated in boldface type and the centimorgan distance between adjacent markers is indicated (small numbers between loci). Markers with the same linkage-map positions are listed together and separated by commas or indicated by branching lines. (B) Magnified version of the linkage map between CBM 9.03 (wx1) and CBM 9.04 (csu147), showing markers around the centromere. The estimated position of the centromere is indicated (solid box, cent9).

RFLP marker loci for which many public probes are available (T. A. Musket and G. L. Davis, University of Missouri-Columbia RFLP Laboratory). Figure 1 shows the genetic map positions of the maize 9 markers

initially chosen for FISH mapping. Very closely linked markers whose order is not resolved are listed together, separated by commas or by branching lines (*e.g.*, sus1, csu43). The estimated position of the centromere (Figure 1, cent9, solid box) resides in bin 9.03 and its region is shown enlarged (Figure 1B). On the basis of this map, the centromere for chromosome 9 was not specifically mapped but was placed between wx1 and csu193 (DAVIS *et al.* 1999).

Selection of sorghum BAC clones for use as FISH probes: We made use of a well-developed fingerprint-contig (FPC) physical map of *S. propinquum* derived from restriction fragment analysis of the YRL BAC library (LIN *et al.* 1999). Clones from this library of partially digested *Hind*III restriction fragments of genomic DNA have an average insert size of 126 kb. A pair of nylon filters containing a gridded array of up to 36,864 YRL BACs are available for hybridization and these filters have been screened with genetic markers such as overgo and RFLP probes (LIN *et al.* 1999; BOWERS *et al.* 2003) and used to select sorghum BACs for use as FISH probes (KOUMBARIS and BASS 2003). High-stringency hybridization ($T_m - 12^\circ$) carried out by us on the YRL sorghum BACs with maize RFLP probes resulted in the detection of an average of 5.3 BACs per probe, consistent with the sixfold genomic coverage of the two filter sets.

This RFLP-based BAC selection procedure is illustrated in Figure 2 for the maize marker, csu145a(pck), and the hybridization results are summarized in Table 1 for all the markers mapped in the study reported here. This marker probe hybridized to 6 BAC clones on the first filter (Figure 2A, left) and 5 on the second (Figure 2A, right). We determined the addresses of these BACs in the library and searched the online Sorghum FPC Map for them (<http://www.genome.arizona.edu/fpc/sorghum/>). Only 8 of the 11 BACs detected by probe csu145 (Figure 2A) were found to be overlapping in a single *S. propinquum* contig, no. 191 (Figure 2D, arrows). BAC a0035C01 was detected by many different maize RFLP probes used and was therefore considered a recurring false positive and excluded from the lists of loci detected. The contig to which the other two BACs (a0028H16 and a0026D19) belong could not be identified with the FPC map. We grew and isolated the remaining 8 detected BACs using a DNA miniprep procedure and subjected them to Southern blot analysis to verify that they did indeed contain a csu145-cross-hybridizing sequence (Figure 2, B and C). The same probe that was used to screen the YRL filters also hybridized to a single band in *Eco*RI-digested DNA minipreps (Figure 2, B and C, lanes 8–15) and to itself (lane 7) but not to other BACs on the same blot (lanes 2–5). Once confirmed by these procedures, a single BAC could be chosen for use as a FISH probe. In some cases, we used a BAC that was previously identified by overgo probe hybridization and subsequently confirmed by us using the corresponding RFLP probe.




FIGURE 2.—Identification, selection, and verification of maize RFLP marker *csu145*-selected sorghum BAC clones. (A) Autoradiographs of *Sorghum propinquum* YRL filters showing six detected clones in the first filter (left) and five in the second (right). BAC a0067L02 is circled and magnified to show one of the unique two-spot patterns from which the BAC identification is decoded. (B and C) An electrophoretic gel and autoradiograph of a blot containing the maize marker *csu145*, the positive control (lane 7), and the eight BACs it detected (lanes 8–15). Also included are the maize marker *csu183* (lane 2), the three clones it detected (lanes 3–5), and the lambda marker (lane 1), which served as negative controls. (D) Fingerprint contig map no. 191 (<http://www.genome.arizona.edu/fpc/sorghum/>) showing the BACs detected by the following markers: five BACs by *cdo1387* (solid circles), seven BACs by *asg44* (asterisks), five BACs by *csu28* (arrowheads), and eight BACs by *csu145* (arrows).

We found that probes for several different but closely linked markers sometimes hybridized to BACs in the same sorghum contig or even to the same sorghum BAC clones (Table 1). The plasmid clones that identified the same contigs but selected different groups of clones are as follows: (1) *csu392* and *csu710* (contig 188); (2) *umc148* and *php10005* (contig 389); and (3) *csu145*, *csu28*, *asg44*, and *cdo1387* (contig 191) (Figure 2D). Sorghum RFLP markers *cdo1387* (Figure 2D, solid circle) and *csu28* (Figure 2D, arrowhead) were already mapped on this physical contig along with other markers and overgo probes. From the positions of the BAC clones detected by maize RFLP markers *asg44* (Figure 2D, asterisks) and *csu145* (Figure 2D, arrows),

we have determined their approximate position on the FPC map (Figure 2D, below the line). From left to right, the order of loci in the contig map for the four mentioned markers is *cdo1387*-*asg44*-*csu28*-*csu145*, the same order as that for the maize linkage map.

Overall, we set out to select BACs for 52 loci to be FISH mapped. We performed BAC filter hybridization for all of the available RFLPs ($n = 47$) and used the information from these or previous overgo hybridizations (BOWERS *et al.* 2003) to select BAC probes for FISH. Nine RFLP probes resulted in a failure to detect sorghum BACs on the YRL BAC grid-array filters ($n = 9$, YN in Table 1) and seven RFLP probes resulted in a failure to detect sorghum BAC restriction fragments at

TABLE 1
Identification and selection of sorghum BACs

Locus	Bin	RFLP size (bp)	Probe type ^c	Probe name	Sorghum FPC contigs	No. of BACs hit	Selected sorghum BACs ^b	FISH probe selected	cM ^c	Mean \pm SEM (<i>n</i>) ^d	FISH locus (cMC) ^e	Cytolocus name
agr118b	9.00	500	R	agr118			YN		0.0			
bn19.07a	9.00	2400	R	bn19.07		7	YN	a0004M18	10.1		9S.79	spb-CBM9.01_S79
umc109 (CBM 9.01) ^f	9.01	800	R	umc109	1220		a0055A15, a0055B21, a0064C07, a0004M18, a0068L03, a0085L22, a0040A04		15.5			(umc109)
umc148	9.01	520	R	umc148	389	5	a0030K10, a0038L23, a0038N20, a0094A07, a0013E22	a0038L23	16.4		FN	
rz144a	9.01	1500	O	SOG1672	389	1	a0030K10	a0030K10	17.4	0.815 \pm 0.004 (15)	9S.82	spb-9.01_S82
rz144c	9.01	1500	O	SOG1672	389	1	a0030K10	a0030K10	21.4	0.745 \pm 0.004 (12)	9S.75	(rz144a) spb-9.01_S75
php10005	9.01	850	R	php10005	389	5	a0073F07, a0014C11, a0068G09, a0045A20, a0017D11	a0045A20	24.8	0.730 \pm 0.002 (31)	9S.73	(rz144c) spb-9.01_S73
csu95a	9.01	1100	O	SOG0620	130	1	a0012H11	a0012H11	27.7	0.678 \pm 0.002 (10)	9S.68	(php10005) spb-9.01_S68
csu95a	9.01	1100	R	csu95	390	8	a0018G19, a0094O05, a0095J06, a0095O12, a0060M11, a0096I01, a0095L23, a0042B21					(csu95a)
umc248a sh1	9.01 9.01	600 2500	R R	umc248 p246A	392	7	YN	a0015M03	27.7		9S.66	spb-9.01_S66
bz1 (CBM 9.02) ^f	9.02	1500	R	pBZ	1308	2	a0084C09, a0094F18, a0021O07, a0015M03, a0088L22, a0021O04, a0091L24	a0015M03	36.4	0.661 \pm 0.001 (18)		(sh1)
csu471	9.02	850	R	csu471	393	11	a0005O05, a0034A09, a0043E13, a0044H06, a0057D08, a0063M03, a0061M22, a0068O17, a0075C01, a0016H09, a0015L15	a0075C01	42.8	0.636 \pm 0.002 (55)	9S.64	spb-CBM9.02_S65
umc256a	9.02	700	R	umc256	805	7	SN	a0020G06	39.3		9S.65	(bz1)
csu486a	9.02	1350	R	csu486	394	8	a0064N21, a0086K22, a0014H21, a0053H23, a0005O15, a0086N18, a0007E04, a0024L14	a0086K22	43.2		9S.39	spb-9.02_S39
prc1	9.02	1080	R	5C02A05	394	9	a0064N21, a0086K22, a0014H21, a0053H23, a0005O15, a0086N18, a0007E04, a0024L14, a0076L04	a0086K22	45.5	0.385 \pm 0.003 (34)		(csu486)
umc105a	9.02	610	R	umc105	396	3	a0016B24, a0084H21, a0054A13	a0076L04	50.2	0.373 \pm 0.002 (42)	9S.37	spb-9.02_S37
csu228 (pfk)	9.02	1190	R	csu228	397	6	a0038I11, a0043G06, a0025D24, a0044C12, a0039P12, a0095E12	a0054A13	54.7	0.276 \pm 0.003 (17)	9S.28	(prc1)
								a0043G06	59.7	0.269 \pm 0.003 (24)	9S.27	spb-9.02_S28
												(umc105)
												spb-9.02_S27
												[csu228 (pfk)]

(continued)

TABLE 1
(Continued)

Locus	Bin	RFLP size (bp)	Probe type ^e	Probe name	Sorghum FPC contigs	No. of BACs hit	Selected sorghum BACs ^b	FISH probe selected	cM ^c	Mean \pm SEM (<i>n</i>) ^d	FISH locus (cMC) ^e	Cytolocus name
wx1 (CBM 9.03)	9.03	2300	G	umc25			131L1 (Bennetzen, UGA)	131L1	63.7	0.133 \pm 0.005 (16)	9S.13	sbb-CBM9.03_S13 (wx1)
wx1 (CBM 9.03)	9.03	2300	R	umc25		3	YN					
tda66d	9.03	300	R	tda66	89	3	a0074N08, a0054I24, a0064A03	a0054I24	63.7	0.031 \pm 0.001 (25)	9S.03	spb-9.03_S03 (tda66d)
cdo17	9.03	900	O	SOG0101	874	5	a0011N04, a0047J04, a0060M19, a0082G20, a0093K24	a0047J04	67.7	0.027 \pm 0.003 (15)	9L.03	spb-9.03_L03 (cdo17)
rf2	9.03	1200	R	rf2a	383	3	a0060G13, a0070K10, a0090D12	a0090D12	67.7		FN	
csu680d	9.03	1550	R	csu680	202	6	a0047C15, a0005O02, a0060L03, a0058B07, a0064H11, a0051F17	a0058B07	67.7		FN	
csu680d	9.03	1550	O	SOG1356	202	3	a0058B07, a0064H11, a0051F17	a0051F17				
bnl7.24a	9.03	2100	R	bnl7.24	1	3	SN					
bnl5.33c	9.03	2100	R	bnl5.33	53	5	a0067P21, a0026E11, a0093B10, a0020L09, a0032H22	a0020L09	68.7	0.043 \pm 0.002 (64)	9L.04	spb-9.03_L04 (bnl5.33c)
csu321	9.03	750	R	csu321	1361	5	a0082L07, a0081N24, a0020F10, a0024F19, a0041D08	a0020F10	69.1	0.035 \pm 0.002 (40)	9L.04	spb-9.03_L04 (csu321)
rgpr3235a	9.03	ND	O	SOG1297	327	1	a0040B08	a0040B08	69.1	0.049 \pm 0.003 (25)	9L.05	spb-9.03_L05 (rgpr3235a)
umc81	9.03	820	R	umc81	14	5	SN		69.6			
csu193	9.03	1270	R	csu193	797	5	a0014C14, a0025K08, a0060B09, a0057H24, a0095I20	a0014C14	70.6		FN	
std6a(dba)	9.03	1200	R	pAS8	797	6	SN		70.6			
gl15	9.03	1150	R	HS1100.43	376	4	a0073J07, a0075D13, a0084N16, a0009B14	a0075D13	70.6	0.063 \pm 0.002 (48)	9L.06	spb-9.03_L06 (gl15)
gl15	9.03	1150	O	SOG0488	376	3	a0073J07, a0075D13, a0084N16					
csu147 (CBM 9.04)	9.04	800	O	SOG1865	69	4	a0022A12, a0049N08, a0054F07, a0068M13	a0049N08	72.8	0.073 \pm 0.003 (21)	9L.07	spb-CBM9.04_L07 (csu147)
csu147 (CBM 9.04)	9.04	800	R	csu147			YN					
csu183b(cdc48)	9.04	850	R	csu183	401	3	SN		78.0			
csu43	9.04	600	R	csu43	550	4	SN		83.1			
sus1	9.04	6000	R	p21.2			YN		83.1			
csu694a(uce)	9.04	950	O	SOG0872	1336	1	a0093O18	a0093O18	84.5	0.355 \pm 0.003 (14)	9L.36	spb-9.04_L36 [csu694a(uce)]
csu694a(uce)	9.04	950	R	csu694	179	6	a0053I22, a0086B07, a0076H16, a0023A07, a0073A14, a0094G22					
umc95 (CBM 9.05)	9.05	660	R	umc95	182	7	a0015A13, a0041F23, a0043K19, a0063J06, a0069D12, a0075C14, a0094N21	a0063J06	86.1	0.377 \pm 0.002 (34)	9L.38	spb-CBM9.05_L38 (umc95)

(continued)

TABLE 1
(Continued)

Locus	Bin	RFLP size (bp)	Probe type ^e	Probe name	Sorghum FPC contigs	No. of BACs hit	Selected sorghum BACs ^b	FISH probe selected	cM ^c	Mean \pm SEM (<i>n</i>) ^d	FISH locus (cMC) ^e	Cytolocus name
csu392a	9.05	1000	O	SOG0184	539	11	a0010M15, a0012B01, a0016C21, a0030A02, a0031D14, a0038B08, a0043J23, a0044E15, a0059K01, a0073M05, a0080L20	a0010M15	97.0	0.413 \pm 0.005 (20)	9L.41	spb-9.05_L41 (csu392a)
csu392a	9.05	1000	O	SOG0184	188	5	a0004C15, a0010J13, a0026E03, a0041K24, a0079J11					
csu392a	9.05	1000	R	csu392	188	6	a0076K14, a0074B22, a0067J12, a0004C15, a0026E03, a0072C14					
csu710e(apx)	9.05	700	O	SOG0484	211	5	a0019J05, a0030K14, a0044G10, a0064D18, a0067N15	a0019J05	101.1	0.456 \pm 0.003 (25)	9L.46	spb-9.05_L46 [csu710e(apx)]
csu710e(apx)	9.05	700	R	csu710	188	5	a0047M11, a0066O15, a0066B02, a0058C12, a0093L24					
csu219(tgd)	9.05	500	R	csu219	189	3	a0059O14, a0071H18, a0067I12	a0059O14	102.3	0.506 \pm 0.006 (43)	9L.51	spb-9.05_L51 [csu219(tgd)]
csu219(tgd)	9.05	500	O	SOG0074	189	2	a0059O14, a0067I12					
csu61a (CBM 9.06)	9.06	500	R	csu61	403	6	a0018G11, a0019H12, a0018J02, a0041I20, a0063F18, a0090O20	a0018G11	104.4		FN	
csu61a (CBM 9.06)	9.06	500	O	SOG1475	403	4	a0018G11, a0019H12, a0018J02, a0041I20					
csu59a	9.06	500	O	SOG0619	270	3	a0019C22, a0074G20, a0084M22	a0074G20	104.4	0.523 \pm 0.002 (24)	9L.52	spb-9.06_L52 (csu59a)
csu59a	9.06	500	R	csu59	189	3	a0059O14, a0071H18, a0067I12	a0071H18				
ibp1	9.06	1150	R	pOD3 ^f			YN		104.4			
csu145a(pck)	9.06	700	R	csu145	191	8	a0043G12, a0041H16, a0046O17, a0052M07, a0067L02, a0068L14, a0079N13, a0093D20	a0093D20	105.4	0.534 \pm 0.003 (26)	9L.53	spb-9.06_L53 [csu145a(pck)]
csu145a(pck)	9.06	700	O	SOG0076	191	11	a0043G12, a0041H16, a0046O17, a0052M07, a0067L02, a0068L14, a0079N13, a0093D20, a0006K22, a0011D09, a0087G20					
csu145a(pck)	9.06	700	O	SOG0076	538	10	a0035K11, a0055A21, a0065B16, a0069H02, a0071B24, a0081F07, a0084G14, a0088L04, a0089F23, a0095K18	a0055A21		0.527 \pm 0.003 (13)	9L.53	spb-9.06_L53 [csu145a(pck)]
dba4	9.06	900	R	pAS14			YN		109.8			
csu28a(rpS22)	9.06	700	R	csu28	191	5	a0010A04, a0032K15, a0076F19, a0087A24, a0093D21	a0093D21	109.8	0.537 \pm 0.003 (50)	9L.54	spb-9.06_L54 [csu28a(rpS22)]
asg44	9.06	500	R	asg44	191	7	a0047G04, a0081A19, a0033A22, a0091J11, a0064A12, a0038E07, a0032I20	a0033A22	110.9	0.624 \pm 0.002 (28)	9L.62	spb-9.06_L62 (asg44)

(continued)

TABLE 1
(Continued)

Locus	RFLP size (bp)	Probe type ^c	Probe name	Sorghum FPC contigs	No. of BACs hit	Selected sorghum BACs ^b	FISH probe selected	cM ^c	Mean \pm SEM (<i>n</i>) ^d	FISH locus (cMC) ^e	Cytolocus name
cdol387(emp70)	9.06	nd	SOG0100	56	4	a0007F13, a0012H21, a0036J08, a0087E20	a0036J08	113.5	0.717 \pm 0.002 (50)	9L.72	spb-9.06.L72
cdol387(emp70)	9.06	nd	SOG0100	191	4	a0016I16, a0036F07, a0036L01, a0093N18	a0036F07		0.722 \pm 0.002 (38)	9L.72	[cdol387(emp70)] spb-9.06.L72
csu1004	9.06	1800	csu1004	1048	5	a0027H23, a0046F16, a0046P22, a0054L18, a0067H10	a0046P22	119.9	0.766 \pm 0.001 (18)	9L.77	[cdol387(emp70)] (csu1004)
asg12 (CBM 9.07)	9.07	700	R asg12	386	3	a0007D16, a0064E21, a0094P12	a0064E21	121.2	0.779 \pm 0.003 (35)	9L.78	spb-CBM9.07.L78 (asg12)
std2a(dba)	9.07	600	R asg12	194	3	a0036L10, a0074H17, a0076L16		122.7			
csu1118	9.07	700	R pAS9	197	1	YN a0078D03	a0078D03	124.3	0.828 \pm 0.003 (25)	9L.83	spb-9.07.L83 (csu1118)
asg59b	9.07	750	R asg59	890	5	SN		132.8			
csu285(his2B)	9.07	800	R csu285	9	8	a0049P09, a0078L14, a0064J04, a0026P15, a0020P12, a0066J18, a0032B18, a0062J05	a0026P15	147.4	0.939 \pm 0.003 (27)	9L.94	spb-9.07.L94 [csu285(his2B)]
csu54b (CBM 9.08) ^f	9.08	1400	R csu54	203	5	a0021O07, a0061D07, a0074A03, a0080B23, a0076H02	a0074A03	150.4		9L.95	spb-CBM9.08.L95 (csu54b)
csu804a(dmp)	9.08	500	R csu804	203	3	a0021O17, a0061D07, a0074A03	a0061D07	150.4		FN	
Xtxa325			G			sbb18256/191b4 (P. E. Klein, TAMU)	sbb18256		0.262 \pm 0.009 (9)	9L.26	sbb_9.04-L26 (Xtxa325)
Xtxp32			G			sbb16685/174g5 (P. E. Klein, TAMU)	sbb16685		0.530 \pm 0.006 (8)	9L.53	sbb_9.06-L53 (Xtxp32)

CBM, core bin marker; TAMU, Texas A&M University.

^a Probe type: R, RFLP; O, overgo probe; G, gift (source is indicated).

^b YN, YRL negative (failure to detect hybridization of RFLP probe on BAC-grid filter arrays); SN, Southern blot negative (failure to detect hybridization of RFLP probe on BAC validation Southern blot).

^c Centimorgan value from the UMC 98 9 linkage map.

^d Numbers represent the mean of the fractional distance of the above-background peak (see MATERIALS AND METHODS) FISH signals on the chromosome arm. SEM, standard error of measurement at 95% confidence level; (*n*) is the number of FISH signals used to calculate the mean.

^e FN, FISH negative, failed to detect FISH signals suitable for mapping on two to three slides.

^f Locus previously mapped by KOUMBARIS and BASS (2003).

^g IBP2 cDNA (from POD3, H. W. BASS, unpublished data).




FIGURE 3.—Image collection and FISH data analysis for the *Waxy1* (*wx1*) locus. (A) The FITC image shows GISH-painted maize pachytene chromosome 9. (B) The rhodamine image shows the signals for *wx1* (arrow). (C) The Cy-5 image shows centromere FISH signals (*cen*). (D) RGB pseudocolor overlay of the DAPI (blue), FITC (red), and rhodamine (green) images. (E) RGB pseudocolor overlay of projections of a straightened chromosome of the FITC (red), rhodamine (green), and Cy-5 (blue) images. FISH-like signals (*) are indicated. (F) Frequency histogram graph for all FISH signals along the short arm of chromosome 9 as described in MATERIALS AND METHODS.

the Southern blot verification step ($n = 7$, SN in Table 1). From a combination of all sources, we obtained a total of 41 BACs that were deemed suitable for FISH mapping as detailed in Table 1.

Pachytene FISH map of maize 9: Pachytene spreads were made from oat–maize 9 anthers. To isolate maize 9 chromosome optically from among the other 21 oat pachytene fibers, we painted the entire maize 9 chromosome by genomic *in situ* hybridization (GISH), using labeled maize total DNA as probe. The FISH method, chromosome straightening, and cytogenetic map position value (centiMcClintock) determination are illustrated in Figure 3; the *S. bicolor* *wx1* BAC 131L1 is used as an example.

All of the FISH experiments involved four-wavelength 3D image collection. Figure 3, A–C, illustrates the three differentially labeled sequences making up the FISH probe cocktail. The DAPI, FITC (maize paint), and rhodamine (BAC) images were overlaid and pseudocolored in blue, red, and green, respectively (Figure 3D). For cytogenetic FISH mapping in maize using pachytene chromosomes, each locus is assigned a centiMcClintock value on the basis of its relative position along the arm as illustrated for the *wx1* locus shown in Figure 3, E and F. Two BAC–FISH signals on the short arm of chromosome 9 were observed in the example shown as green spots (Figure 3E, asterisks). We typically straightened >40 chromosomes per locus, and in this case we detected a total of 53 discrete signals on maize 9S. To distinguish

true signal from background, we visualized the distribution of signals as a frequency histogram (Figure 3F). The average location of the above-background FISH signals was determined to be at 9S.13. Thus the locus is designated as *sbb-CBM9.03_S13 (wx1)*, indicating that an *S. bicolor* BAC clone was used as a FISH probe for the *wx1* locus that is core bin marker 9.03 and that the resulting locus is at centiMcClintock position 9S.13.

From a total of 41 BACs that were used as FISH probes, 36 of them yielded suitable FISH mapping data (Table 1). Representative images for 24 newly mapped loci are shown in Figure 4. These illustrate FISH probes distributed across the entire chromosome, including 8 from the short arm (Figure 4, A–H) and 16 from the long arm (Figure 4, I–X). Each section in Figure 4 corresponds to a single FISH probe and shows a representative straightened-arm image, the frequency histogram of FISH signals (as described for Figure 3), and the cytogenetic FISH locus name. All of the probes mapped to a single locus except for the *rz144*-selected BAC (a0030K10) that hybridized to two closely linked regions, 9S.75 and 9S.82 (Figure 4A).

The overgo and RFLP probes did not always detect the exact same set of BACs, as exemplified with *csu145* (Table 1). In this case, we attempted FISH with two different BACs, the RFLP-selected a0093D20 and the overgo-selected a0055A21, both of which were FISH mapped to 9L.53. In a few other cases, very tightly linked maize markers even hybridized to the same BACs. For




FIGURE 4.—FISH signal histograms and locus names for 24 BAC-FISH probes. Each section shows the histogram, the probe name, and the representative arm projection as described in Figure 3, E and F. The sorghum BAC name is indicated in brackets for each section. The loci are located on the short (A–H) or the long (I–X) arm of maize pachytene chromosome 9.




FIGURE 5.—Refining the linkage map position of cent9, the maize 9 centromere. Integrated linkage (A) and cytogenetic (B) maps showing the new estimated map position (A, hatched box) for the maize 9 centromere are shown. Map units for A are described in the Figure 1 legend. Map units for B are centiMcClintocks, and 9S refers to the short arm and 9L refers to the long arm. (C) Straightened projections of chromosomes showing the representative of each of the centromere-linked markers, pseudocolored as described in Figure 3E.

instance, the markers *csu486a* and *prc1* independently detected eight common BACs in contig 389 (Table 1). However, one extra BAC (*a0076L04*) was detected only by *prc1*, which was then used as a FISH probe in addition to BAC *a0086K22* that represented *csu486a*. Using these BACs, we FISH mapped *prc1* to 9S.37 while *csu486a* mapped to 9S.39 (Table 1). Three other locus pairs (*umc148* and *rz144*, *csu219* and *csu59a*, and *csu804a* and *csu54b*) showed this same sort of BAC codetection result. The distances separating these locus pairs range from 0 to 5 cM, indicating that this spacing may be near the axial resolution limit for FISH mapping in some regions.

Cytological evidence for regions of genome hyperexpansion: Three distinct noncentromeric regions of the chromosome showed evidence of hyperexpansion relative to the overall increase of 3.3-fold for maize *vs.* sorghum genome size. These regions are delimited by three pairs of loci, one on 9S and two on 9L. To compare them, we used the following map distance values for maize chromosome 9: 9S is 67 cM, 100 cMC, and 77 Mbp long and 9L is 84 cM, 100 cMC, and 114 Mbp long (BENNETT and LAURIE 1995; DAVIS *et al.* 1999; ANDERSON *et al.* 2004).

The first of these three hyperexpanded regions is in bin 9.02 on 9S between *csu471* and *csu486a* at 9S.64–9S.39. In this segment of the chromosome, the cM/cMC ratio is 0.11, 84% lower than the whole-arm average. The second such region is in bin 9.04 on 9L between *csu147* and *csu694a(uce)* at 9L.07–9L.36. In this segment of the chromosome, the cM/cMC ratio is 0.40, 52% lower than the whole-arm average. The third case is in bin 9.06 on 9L between *csu28a(rpS22)* and *cdo1387a(emp70)* at 9L.54–9L.72, a segment with a length of 3.7 cM and 18 cMC. The cM/cMC ratio in this region is 0.21, 76% lower than the whole-arm average. In other words, these three regions

show sixfold, twofold, and fourfold higher amounts of chromosomal distance than expected from the whole-arm average ratios of 0.67 (short) and 0.84 (long) cM/cMC.

In this third region [*csu28a(rpS22)* to *cdo1387a(emp70)* in maize bin 9.06], the corresponding sorghum markers are found in a single well-defined sorghum contig (no. 191). This allows for a comparison of the genome expansion rates for this region relative to the overall expansion rate of maize relative to sorghum. These markers in sorghum are separated by ~200 Kbp, whereas they are separated in the cytogenetic map by 18 cMC. Using the estimate of 1.14 Mbp/cMC for 9L, we calculate that this region has expanded at a rate 25-fold higher than that of the overall maize genome relative to the sorghum genome. This is one of the three major regions that were found to exhibit this unexpected pattern of marker distribution, which we refer to here as regions of genome hyperexpansion.

FISH mapping of CentC within centromeric loci: The positions of centromeres on linkage maps are generally not well defined because they typically lack alleles or polymorphisms suitable for standard linkage mapping. For example, the location of the maize 9 centromere is an estimate that spans a large amount of bin 9.03. To map the maize 9 centromere more precisely relative to closely linked loci, we mapped the centromere-linked markers *wx1*, *tda66d*, *cdo17*, *bnl5.33c*, *csu321*, *rgpr3235a*, *gl15*, and *csu147* as shown in Figure 5. Representative FISH images of straightened pachytene chromosomes (Figure 5C) are shown and the location of these loci on the cytogenetic map is indicated (Figure 5B). We mapped *wx1* and *tda66d* in the short arm, whereas the other markers were in the long arm (Figure 5B). From these data, we refined the estimated location of the centromere to a smaller region




FIGURE 6.—Integrated maps of maize chromosome 9. (A) Fingerprint contig numbers for the Sorghum FPC map and their associated linkage groups. (B) The maize 9 linkage map is shown as described in Figure 1, except that the cent9 location reflects the revised position as described in Figure 5. The names of the sorghum BACs used as a FISH probe are listed after the slashes (after the slashes). (C) The new cytogenetic FISH 9 map showing the location (solid double circles) of loci along the short (9S) and the long (9L) arm. The *S. bicolor* BAC sbb18256 (Xtxa325) at 9L.27 is not located on the linkage map and the *S. bicolor* BAC sbb16685 (Xtxp32) was mapped to 9L.53, but is not shown. The new cytogenetic loci names are displayed at the right of the map in C and are detailed in Table 1.

(Figure 5A, hatched box), between tda66d (9S.03) and cdo17 (9L.03) (Figure 5B). This approach allows for assignment of a given locus to either the long arm or the short arm and provides an indirect method for delimiting the location of the centromere on linkage maps, assuming that the CentC repeat cluster faithfully reports the position of the functional centromere.

Integrating data from sorghum and maize maps: The FISH-BAC probes used in this study are unique in their applicability to multiple maps in maize and sorghum. The new cytogenetic FISH map is summarized in Figure 6 and the way in which it was created provides direct links between the physical map of sorghum (Figure 6A) and the linkage map of maize (Figure 6B). The results

anchor sorghum BACs to maize pachytene chromosomes by high-stringency DNA sequence hybridization. We found nearly complete colinearity between maize linkage and cytogenetic maps, but the distributions and relative spacing between markers did not always match. One observed partial violation of agreement came from data from the sorghum BAC for rz144. In this case, rz144a and rz144c are both located proximal to umc109 on the UMC 98 9 linkage map, but the FISH signals were found on opposite sides of umc109 on the cytogenetic map. This order of loci from the telomeric end (rz144a-umc109-rz144c) is the same as that of a newer linkage map of maize, the IBM map. From these overall findings, we can conclude that this method is robust and

suitable for development of a detailed cytogenetic FISH map of maize.

DISCUSSION

Cytogenetic mapping provides information on the structure and evolution of genomes (KORENBERG *et al.* 1999; CHEUNG *et al.* 2001; TRASK 2002; GONZALEZ *et al.* 2005). In addition, the BAC–FISH approach can be combined with bioinformatics to resolve inconsistencies among genome-mapping data or reveal ancestral relationships between closely related species (MA *et al.* 2006; reviewed by ROCCHI *et al.* 2006). In maize, the use of molecular cytogenetics has great potential for shedding light on the well-documented structural diversity of its chromosomes and for contributing to ongoing genome-assembly efforts (MARTIENSSSEN *et al.* 2004; MESSING and DOONER 2006). Furthermore, the development of cytogenetic resources for maize will facilitate investigations regarding the origin and evolution of the maize genome and its relationship to that of its close relatives, such as sorghum (GAUT and DOEBLEY 1997; GAUT 2001; KATO *et al.* 2004, 2005; SWIGONOVA *et al.* 2004; BOWERS *et al.* 2005; HABERER *et al.* 2005; BRUGGMANN *et al.* 2006; LAMB and BIRCHLER 2006; SHERIDAN and AUGER 2006).

The development of linkage and FPC-type physical maps of maize has led to a good understanding of the distribution of genes along the chromosomes (DAVIS *et al.* 1999; LEE *et al.* 2002; SHAROPOVA *et al.* 2002) and the physical distances between genes that reside within single FPC contigs (NELSON *et al.* 2005; PAMPANWAR *et al.* 2005; YIM *et al.* 2007). Even so, the linkage maps are based on recombination frequencies that vary widely in relation to physical distances (ANDERSON *et al.* 2003, 2004; WANG *et al.* 2006), and the FPC physical maps may not accurately predict the physical distance between the ends of adjacent contigs. Even as the maize genome sequence approaches completion, cytogenetic tools will remain useful for evaluating the variation between different species, subspecies, and cultivars within the genus *Zea* (WHITE and DOEBLEY 1998; LIU *et al.* 2003; KATO *et al.* 2004; BUCKLER *et al.* 2006; LAMB and BIRCHLER 2006; LAMB *et al.* 2007).

The pachytene FISH map of maize 9: Here we have described the physical organization of maize chromosome 9 by creating a new cytogenetic BAC–FISH map that integrates maize maps, provides links to sorghum maps, and uncovers several hotspots of maize genome expansion that were not anticipated from prior maps or comparative sequence analysis (DAVIS *et al.* 1999). The strategy of cross-species mapping using conserved gene sequences has proven successful in examining genome structures and relationships and predicting locations of genes and DNA markers in related species (HULBERT *et al.* 1990; FUCHS *et al.* 1996; GOMEZ *et al.* 1997; GALE and DEVOS 1998; ZWICK *et al.* 1998; JACKSON *et al.* 2000;

DRAYE *et al.* 2001; KOUMBARIS and BASS 2003; DEVOS 2005). In this study, >40 sorghum BACs were used as FISH probes to create 32 new cytogenetic loci, most of which correspond to RFLP loci that are well mapped and widely used in maize linkage analysis. The success rate of this approach establishes it as a valuable and informative method for developing a detailed, integrated cytogenetic FISH map of maize. Starting with the maize 9 linkage map, 52 markers were selected and 9 of them failed at the BAC filter screening, 7 of them failed at the Southern blot stage, and 6 of them failed at the FISH stage (Table 1).

The markers were nearly completely colinear between the genetic and cytogenetic maps, but irregularly distributed along the entire length of the chromosome. The sort of differences between linkage and physical distances along individual chromosomes that we found is well documented for maize, sorghum, and other plant species (GILL *et al.* 1996; PETERSON *et al.* 1999; KUNZEL *et al.* 2000; SADDER and WEBER 2002; KIM *et al.* 2005a; WANG *et al.* 2006). For example, the centromere and other heterochromatic regions often exhibit significantly fewer linkage-map units per unit physical distance because they typically have relatively low rates of meiotic recombination (SHERMAN and STACK 1995; PETERSON *et al.* 1999; HARPER and CANDE 2000; ANDERSON *et al.* 2003, 2004).

Local variation in marker spacing: To compare the different maps of the maize and sorghum genomes, we calculated the overall ratio of DNA length in physical units (base pairs or centiMcClintocks) to the genetic length in map units (centimorgans). This overall ratio is calculated separately for each arm because each arm encompasses 100 cMC (LAWRENCE *et al.* 2006). The DNA content for the arms of maize chromosomes in line Seneca 60, the pollen parent for oat–maize 9, was determined by BENNETT and LAURIE (1995) to be 77 Mbp for 9S and 114 Mbp for 9L. The numbers of linkage-map units from the UMC 98-based “Genetic 2005 9” linkage map are 67 cM for 9S and 84 cM for 9L (<http://www.maizegdb.org/cgi-bin/displaymaprecord.cgi?id=940888>). From this information, we derive the values per centiMcClintock to be 0.77 Mbp/cMC for 9S, 1.14 Mbp/cMC for 9L, 0.67 cM/cMC for 9S, and 0.84 cM/cMC for 9L.

Large deviations in the cM/cMC ratios were observed when two intervals on 9S were compared: the *csu95a*–*sh1* region (9S.68–9S.66) in bin 9.01 and the *csu228*(*pfk*)–*wx1* region (9S.27–9S.13) in bin 9.03. The distance between *csu95a* and *sh1* is 8.7 cM but corresponding to only 2.0 cMC, whereas the distance between *csu228*(*pfk*) and *wx1* is 4.0 cM, corresponding to 14.0 cMC. The cM/cMC ratios for these two regions are 4.35 and 0.286, respectively, representing a 6.5-fold increase and a 57% decrease relative to the cM/cMC ratio averaged over the whole arm of 9S.

In addition to these variations in the frequency of recombination per cytological distance, additional analysis

of marker distribution uncovered striking evidence of regions where the maize genome appears to exhibit accelerated expansion. One of these regions is the segment on 9L between *csu28a*(*rpS22*) and *cdo1387a*(*emp70*) in bin 9.06. These markers are only 3.7 cM apart, but separated by 18 cMC on the basis of our FISH map. This region exhibits a 25-fold higher genome expansion rate than that of the whole-genome average for maize relative to sorghum. In another region, between *csu471* and *csu486a* on 9S in bin 9.02, we examined the physical map covered by maize FPC contigs 371, 372, and 373 (<http://www.maizesequence.org>). The distal marker (*csu471*) is anchored in maize contig 371, but the proximal marker (*csu486a*) is not anchored and may be in contig 372 or 373. Although this segment is only 2.7 cM in length, it spans 25 cMC on the basis of our FISH map. This physical distance is predicted to represent ~19.25 Mbp, suggesting that the gap between maize contigs 371 and 372, or between 372 and 373, may be unusually large compared to other contig gaps. Findings such as these may be important for guiding positional cloning or other strategies that rely on chromosome walking in this area. They may also reveal hotspots for insertions or duplications of mobile or other repetitive sequence elements. It will be of interest to examine the gene content and arrangement of genes in these regions.

In considering possible mechanisms for this hyper-expansion, this region does not appear to correspond to any known heterochromatic knobs or other large blocks of gene-depleted chromatin that could account for this variation. Determining whether these regions are common to other lines of maize or unique to the pollen parent line, Seneca 60, will be of interest. Recent analysis of the maize genome has also revealed uneven chromosome contraction and expansion, but on a slightly smaller scale (BRUGGMANN *et al.* 2006). BRUGGMANN *et al.* (2006) attributed the localized chromosome expansion to the insertion of retrotransposable elements. An alternative explanation is that these expanded areas are peculiar to the oat-maize addition lines and result from chromosome rearrangements or amplifications that could have occurred after the oat-maize cross. This idea, however, is not consistent with observations from oat-maize 6 lines in which the maize 6 centromere region was found to be structurally stable in three independent addition lines (JIN *et al.* 2004). Comparative analysis of these regions in related species may provide insight into their origin and significance.

Concordance of cytogenetic mapping data from maize: The linear order of markers in the cytogenetic FISH map was fully concordant with that of the linkage map except for *rz144* and a very minor switch between two very tightly linked loci (*bnl.5.33c* and *csu321*). In the UMC 98/Genetic 2005 map of chromosome 9, the locus order is *umc109*, *rz144a*, *rz144c*, whereas their order in the IBM2 2004 neighbors 9 map is *rz144a*, *umc109*,

rz144c. The sorghum BAC-FISH probe *a0030K10* hybridized to two loci, one at 9S.82 and one at 9S.75. These two loci flank the position determined for the FISH signal for *umc109*. The only locus-order inconsistency between the cytogenetic FISH map and the UMC 98 linkage map also occurred between the two linkage maps; the FISH data matched the IBM2 map data.

Map discrepancies in maize are not uncommon because of the inherent differences in genomic structure between different lines of maize (FU and DOONER 2002; LEE *et al.* 2002; BRUGGMANN *et al.* 2006). Discrepancies can also result from errors due to small sample sizes or from differences in mapping techniques. For example, the *Waxy1* locus is particularly variable among different maps, having been mapped to 9S and 9L in linkage studies and to different regions on 9S in cytogenetic studies. The *Waxy1* locus has been localized to ~9S.02–9S.06 by SHEN *et al.* (1987), 9S.06 by ANDERSON *et al.* (2004), 9S.46 by WANG *et al.* (2006), and 9S.13 by us. All these studies were based on pachytene chromosomes, but ANDERSON *et al.* (2004) extrapolated the position from inbred KYS recombination nodule distribution, WANG *et al.* (2006) used direct FISH on KYS chromosomes that were uniformly elongated by extra pepsin treatment, and the position in our study was from a sorghum BAC-FISH probe hybridized to Seneca 60 chromosomes carried in an oat-genome background. The data, thus, may not be directly comparable. The sorghum BAC probe for *wx1* has been sequenced and found to be syntenic with the maize and rice *waxy1* region (J. MA and J. L. BENNETZEN, personal communication). To date, the cytological maps are in good agreement with each other and the linkage maps in terms of the linear order of the loci. The map positions we detected were self-consistent, and the relative positions did not vary with absolute length of the pachytene-stage chromosomes themselves. Development of cytogenetic FISH maps for maize (SADDER and WEBER 2001; CHENG *et al.* 2002; KOUMBARIS and BASS 2003; ANDERSON *et al.* 2004; SHERIDAN and AUGER 2006; WANG *et al.* 2006) may eventually comprise multiple maps generated by various techniques for different genotypes.

Centromere mapping: FISH mapping of BAC clones around the centromere provided increased resolution for extrapolating the position of the maize 9 centromere within the linkage maps. We were able to map eight markers on this region using carefully chosen sorghum BAC clones. By coincident staining of the CentC repeats and the BAC-FISH probes on individual fibers, we could make chromosome-arm assignment around the centromere. The centromere, as marked by the CentC cluster, is located between *tda66d* at 9S.03 and *cdo17* at 9L.03. This position is close to but not the same as that determined by LUCE *et al.* (2006), using a CenH3-based method of centromere mapping. Placing centromeres on the genetic map is an indirect form of

linkage mapping and it is further complicated by the dynamic nature of centromeres and their epigenetic specification (NAGAKI *et al.* 2004; LEE *et al.* 2005; LUCE *et al.* 2006).

Mapping loci for markers <1 kbp in length: One of the major challenges for cytogenetic mapping is sensitivity, with the goal of specifically detecting small segments of DNA. RFLP probes have been used in thousands of linkage studies in maize since they were first developed >20 years ago. Localizing these markers cytogenetically would therefore be a valuable extension of maize genetics resources. For example, the RFLP probe asg44 is 500 bp, and probe csu145 is 700 bp. In the study reported here, the advance of KOUMBARIS and BASS (2003) was employed to map >30 loci with an average size near 1 kbp. To date, no robust technology is available for FISH mapping of DNA sequences of this size, but use of RFLP-selected sorghum BACs as FISH probes allowed us to bypass this limitation while adding a new dimension, direct links to the sorghum genome, to the resulting cytological data.

In summary, we have produced a high-resolution FISH map of maize pachytene chromosome 9 with BAC probes from sorghum genomic DNA. We found that the loci were mostly colinear between the linkage and the cytological maps of maize and that regions of genome hyperexpansion could be detected by comparative analysis of various maps containing shared markers. These findings serve to integrate genetic data across different maize maps. They also serve to generate new links between the maps of the maize and sorghum genomes. Our results have important implications for understanding and exploring the structure and the evolution of the maize genome while developing new reagents for chromosome research in the grasses.

We thank Debbie Figueroa and Shaun Murphy for critical reading and helpful comments on the manuscript. We thank R. J. (Bobbye) Hill for her support and assistance with this project. We thank P. E. Klein (Texas A&M University) and J. L. Bennetzen (University of Georgia) for providing some of the *S. bicolor* BACs used in this study. This work was supported by the National Science Foundation (DBI-0321639).

LITERATURE CITED

- ADAWY, S. S., R. M. STUPAR and J. JIANG, 2004 Fluorescence in situ hybridization analysis reveals multiple loci of knob-associated DNA elements in one-knob and knobless maize lines. *J. Histochem. Cytochem.* **52**: 1113–1116.
- ANANIEV, E. V., R. L. PHILLIPS and H. W. RINES, 1998 Chromosome-specific molecular organization of maize (*Zea mays* L.) centromeric regions. *Proc. Natl. Acad. Sci. USA* **95**: 13073–13078.
- ANDERSON, L. K., G. G. DOYLE, B. BRIGHAM, J. CARTER, K. D. HOOKER *et al.*, 2003 High-resolution crossover maps for each bivalent of *Zea mays* using recombination nodules. *Genetics* **165**: 849–865.
- ANDERSON, L. K., N. SALAMEH, H. W. BASS, L. C. HARPER, W. Z. CANDE *et al.*, 2004 Integrating genetic linkage maps with pachytene chromosome structure in maize. *Genetics* **166**: 1923–1933.
- BENNETT, M. D., and D. A. LAURIE, 1995 Chromosome size in maize and sorghum using em serial section reconstructed nuclei. *Maydica* **40**: 199–204.
- BENNETZEN, J. L., V. L. CHANDLER and P. SCHNABLE, 2001 National Science Foundation-sponsored workshop report. Maize genome sequencing project. *Plant Physiol.* **127**: 1572–1578.
- BOWERS, J. E., C. ABBEY, S. ANDERSON, C. CHANG, X. DRAYE *et al.*, 2003 A high-density genetic recombination map of sequence-tagged sites for sorghum, as a framework for comparative structural and evolutionary genomics of tropical grains and grasses. *Genetics* **165**: 367–386.
- BOWERS, J. E., M. A. ARIAS, R. ASHER, J. A. AVISE, R. T. BALL *et al.*, 2005 Comparative physical mapping links conservation of microsynteny to chromosome structure and recombination in grasses. *Proc. Natl. Acad. Sci. USA* **102**: 13206–13211.
- BRUGGMANN, R., A. K. BHARTI, H. GUNDLACH, J. LAI, S. YOUNG *et al.*, 2006 Uneven chromosome contraction and expansion in the maize genome. *Genome Res.* **16**: 1241–1251.
- BUCKLER, E. S., B. S. GAUT and M. D. MCMULLEN, 2006 Molecular and functional diversity of maize. *Curr. Opin. Plant Biol.* **9**: 172–176.
- BURR, B., F. A. BURR, K. H. THOMPSON, M. C. ALBERTSON and C. W. STUBER, 1988 Gene mapping with recombinant inbreds in maize. *Genetics* **118**: 519–526.
- CARLSON, W. R., 1988 The cytogenetics of corn, pp. 259–344 in *Corn and Corn Improvement*, edited by G. F. SPRAGUE and J. W. DUDLEY. American Society of Agronomy, Madison, WI.
- CAUSSE, M., S. SANTONI, C. DAMERVAL, A. MAURICE, A. CHARCOSSET *et al.*, 1996 A composite map of expressed sequences in maize. *Genome* **39**: 418–432.
- CHANDLER, V. L., and V. BRENDEL, 2002 The Maize Genome Sequencing Project. *Plant Physiol.* **130**: 1594–1597.
- CHEN, C. C., C. M. CHEN, F. C. HSU, C. J. WANG, J. T. YANG *et al.*, 2000 The pachytene chromosomes of maize as revealed by fluorescence in situ hybridization with repetitive DNA sequences. *Theor. Appl. Genet.* **101**: 30–36.
- CHENG, Z., C. R. BUELL, R. A. WING, M. GU and J. JIANG, 2001a Toward a cytological characterization of the rice genome. *Genome Res.* **11**: 2133–2141.
- CHENG, Z., G. G. PRESTING, C. R. BUELL, R. A. WING and J. JIANG, 2001b High-resolution pachytene chromosome mapping of bacterial artificial chromosomes anchored by genetic markers reveals the centromere location and the distribution of genetic recombination along chromosome 10 of rice. *Genetics* **157**: 1749–1757.
- CHENG, Z., C. R. BUELL, R. A. WING and J. JIANG, 2002 Resolution of fluorescence in-situ hybridization mapping on rice mitotic prometaphase chromosomes, meiotic pachytene chromosomes and extended DNA fibers. *Chromosome Res.* **10**: 379–387.
- CHEUNG, V. G., N. NOWAK, W. JANG, I. R. KIRSCH, S. ZHAO *et al.*, 2001 Integration of cytogenetic landmarks into the draft sequence of the human genome. *Nature* **409**: 953–958.
- COE, E. H., 1994 A-a translocations: breakpoints and stocks., pp. 364–376 in *The Maize Handbook*, edited by M. FREELING and V. WALBOT. Springer-Verlag, New York.
- COE, E., D. HOISINGTON and M. NEUFFER, 1987 Linkage map of corn (maize) (*Zea mays* L.). *Maize Genet. Crop. Newslett.* **61**: 116–147.
- CREIGHTON, H. B., and B. MCCLINTOCK, 1931 A correlation of cytological and genetical crossing over in *Zea mays*. *Proc. Natl. Acad. Sci. USA* **17**: 492–497.
- DAVIS, G. L., M. D. MCMULLEN, C. BAYSDORFER, T. MUSKET, D. GRANT *et al.*, 1999 A maize map standard with sequenced core markers, grass genome reference points and 932 expressed sequence tagged sites (ESTs) in a 1736-locus map. *Genetics* **152**: 1137–1172.
- DE JONG, J. H., P. FRANSZ and P. ZABEL, 1999 High resolution FISH in plants - techniques and applications. *Trends Plant Sci.* **4**: 258–263.
- DEMPSEY, E., 1994 Traditional analysis of maize pachytene chromosomes., pp. 432–441 in *The Maize Handbook*, edited by M. FREELING and V. WALBOT. Springer-Verlag, New York.
- DESEL, C., C. JUNG, D. CAI, M. KLEINE and T. SCHMIDT, 2001 High-resolution mapping of YACs and the single-copy gene Hs1 (pro-1) on *Beta vulgaris* chromosomes by multi-colour fluorescence in situ hybridization. *Plant Mol. Biol.* **45**: 113–122.
- DEVOS, K. M., 2005 Updating the 'crop circle'. *Curr. Opin. Plant Biol.* **8**: 155–162.
- DONG, F., J. SONG, S. NAESS, J. HELGESON, C. GEBHARDT *et al.*, 2000 Development and applications of a set of chromosome-specific cytogenetic DNA markers in potato. *Theor. Appl. Genet.* **101**: 1001–1007.

- DRAYE, X., Y. R. LIN, X. Y. QIAN, J. E. BOWERS, G. B. BUROW *et al.*, 2001 Toward integration of comparative genetic, physical, diversity, and cytomolecular maps for grasses and grains, using the sorghum genome as a foundation. *Plant Physiol.* **125**: 1325–1341.
- EMERSON, R., G. BEADLE and A. FRASER, 1935 A summary of linkage studies in maize. *Cornell Univ. Agric. Exp. Stn. Memoir* **180**: 1–83.
- FRANSZ, P. F., C. ALONSO-BLANCO, T. B. LIHARSKA, A. J. PEETERS, P. ZABEL *et al.*, 1996 High-resolution physical mapping in *Arabidopsis thaliana* and tomato by fluorescence *in situ* hybridization to extended DNA fibres. *Plant J.* **9**: 421–430.
- FU, H., and H. K. DOONER, 2002 Intrasppecific violation of genetic colinearity and its implications in maize. *Proc. Natl. Acad. Sci. USA* **99**: 9573–9578.
- FU, Y., T. J. WEN, Y. I. RONIN, H. D. CHEN, L. GUO *et al.*, 2006 Genetic dissection of intermated recombinant inbred lines using a new genetic map of maize. *Genetics* **174**: 1671–1683.
- FUCHS, J., A. HOUBEN, A. BRANDES and I. SCHUBERT, 1996 Chromosome 'painting' in plants—A feasible technique? *Chromosoma* **104**: 315–320.
- GALE, M. D., and K. M. DEVOS, 1998 Comparative genetics in the grasses. *Proc. Natl. Acad. Sci. USA* **95**: 1971–1974.
- GARDINER, J., S. SCHROEDER, M. L. POLACCO, H. SANCHEZ-VILLEDA, Z. FANG *et al.*, 2004 Anchoring 9,371 maize expressed sequence tagged unigenes to the bacterial artificial chromosome contig map by two-dimensional overgo hybridization. *Plant Physiol.* **134**: 1317–1326.
- GAUT, B. S., 2001 Patterns of chromosomal duplication in maize and their implications for comparative maps of the grasses. *Genome Res.* **11**: 55–66.
- GAUT, B. S., and J. F. DOEBLEY, 1997 DNA sequence evidence for the segmental allotetraploid origin of maize. *Proc. Natl. Acad. Sci. USA* **94**: 6809–6814.
- GAUT, B. S., M. LE THIERRY D'ENNEQUIN, A. S. PEEK and M. C. SAWKINS, 2000 Maize as a model for the evolution of plant nuclear genomes. *Proc. Natl. Acad. Sci. USA* **97**: 7008–7015.
- GILL, K. S., B. S. GILL, T. R. ENDO and E. V. BOYKO, 1996 Identification and high-density mapping of gene-rich regions in chromosome group 5 of wheat. *Genetics* **143**: 1001–1012.
- GOMEZ, M., M. NURUL ISLAM-FARIDI, S.-S. WOO, K. F. SCHERTZ, D. CZESCHIN *et al.*, 1997 FISH of a maize sh2-selected sorghum BAC to chromosomes of *Sorghum bicolor*. *Genome* **40**: 475–478.
- GONZALEZ, J., M. NEFEDOV, I. BOSDET, F. CASALS, O. CALVETE *et al.*, 2005 A BAC-based physical map of the *Drosophila buzzatii* genome. *Genome Res.* **15**: 885–892.
- HABERER, G., S. YOUNG, A. K. BHARTI, H. GUNDLACH, C. RAYMOND *et al.*, 2005 Structure and architecture of the maize genome. *Plant Physiol.* **139**: 1612–1624.
- HANSON, R. E., M. S. ZWICK, S. CHOI, M. N. ISLAM-FARIDI, T. D. MCKNIGHT *et al.*, 1995 Fluorescent *in situ* hybridization of a bacterial artificial chromosome. *Genome* **38**: 646–651.
- HARPER, L. C., and W. Z. CANDE, 2000 Mapping a new frontier; development of integrated cytogenetic maps in plants. *Funct. Integr. Genomics* **1**: 89–98.
- HARUSHIMA, Y., M. YANO, A. SHOMURA, M. SATO, T. SHIMANO *et al.*, 1998 A high-density rice genetic linkage map with 2275 markers using a single F2 population. *Genetics* **148**: 479–494.
- HASS-JACOBUS, B. L., M. FUTRELL-GRIGGS, B. ABERNATHY, R. WESTERMAN, J. L. GOICOECHEA *et al.*, 2006 Integration of hybridization-based markers (overgos) into physical maps for comparative and evolutionary explorations in the genus *Oryza* and in *Sorghum*. *BMC Genomics* **7**: 199.
- HELENTJARIS, T., D. F. WEBER and S. WRIGHT, 1986 Use of monosomics to map cloned DNA fragments in maize. *Proc. Natl. Acad. Sci. USA* **83**: 6035–6039.
- HOWELL, E. C., G. C. BARKER, G. H. JONES, M. J. KEARSEY, G. J. KING *et al.*, 2002 Integration of the cytogenetic and genetic linkage maps of *Brassica oleracea*. *Genetics* **161**: 1225–1234.
- HULBERT, S. H., T. E. RICHTER, J. D. AXTELL and J. L. BENNETZEN, 1990 Genetic mapping and characterization of sorghum and related crops by means of maize DNA probes. *Proc. Natl. Acad. Sci. USA* **87**: 4251–4255.
- ISLAM-FARIDI, M. N., K. L. CHILDS, P. E. KLEIN, G. HODNETT, M. A. MENZ *et al.*, 2002 A molecular cytogenetic map of sorghum chromosome 1: fluorescence *in situ* hybridization analysis with mapped bacterial artificial chromosomes. *Genetics* **161**: 345–353.
- JACKSON, S. A., Z. CHENG, M. L. WANG, H. M. GOODMAN and J. JIANG, 2000 Comparative fluorescence *in situ* hybridization mapping of a 431-kb *Arabidopsis thaliana* bacterial artificial chromosome contig reveals the role of chromosomal duplications in the expansion of the *Brassica rapa* genome. *Genetics* **156**: 833–838.
- JIANG, J., B. S. GILL, G. L. WANG, P. C. RONALD and D. C. WARD, 1995 Metaphase and interphase fluorescence *in situ* hybridization mapping of the rice genome with bacterial artificial chromosomes. *Proc. Natl. Acad. Sci. USA* **92**: 4487–4491.
- JIN, W., J. R. MELO, K. NAGAKI, P. B. TALBERT, S. HENIKOFF *et al.*, 2004 Maize centromeres: organization and functional adaptation in the genetic background of oat. *Plant Cell* **16**: 571–581.
- KATO, A., J. C. LAMB and J. A. BIRCHLER, 2004 Chromosome painting using repetitive DNA sequences as probes for somatic chromosome identification in maize. *Proc. Natl. Acad. Sci. USA* **101**: 13554–13559.
- KATO, A., J. M. VEGA, F. HAN, J. C. LAMB and J. A. BIRCHLER, 2005 Advances in plant chromosome identification and cytogenetic techniques. *Curr. Opin. Plant Biol.* **8**: 148–154.
- KIM, J. S., M. N. ISLAM-FARIDI, P. E. KLEIN, D. M. STELLY, H. J. PRICE *et al.*, 2005a Comprehensive molecular cytogenetic analysis of sorghum genome architecture: distribution of euchromatin, heterochromatin, genes and recombination in comparison to rice. *Genetics* **171**: 1963–1976.
- KIM, J. S., P. E. KLEIN, R. R. KLEIN, H. J. PRICE, J. E. MULLETT *et al.*, 2005b Molecular cytogenetic maps of sorghum linkage groups 2 and 8. *Genetics* **169**: 955–965.
- KORENBERG, J. R., X. N. CHEN, Z. SUN, Z. Y. SHI, S. MA *et al.*, 1999 Human genome anatomy: BACs integrating the genetic and cytogenetic maps for bridging genome and biomedicine. *Genome Res.* **9**: 994–1001.
- KOUMBARIS, G. L., and H. W. BASS, 2003 A new single-locus cytogenetic mapping system for maize (*Zea mays* L.): overcoming FISH detection limits with marker-selected sorghum (*S. propinquum* L.) BAC clones. *Plant J.* **35**: 647–659.
- KULIKOVA, O., G. GUALTIERI, R. GEURTS, D. J. KIM, D. COOK *et al.*, 2001 Integration of the FISH pachytene and genetic maps of *Medicago truncatula*. *Plant J.* **27**: 49–58.
- KUMAR, A., and J. L. BENNETZEN, 1999 Plant retrotransposons. *Annu. Rev. Genet.* **33**: 479–532.
- KUNZEL, G., L. KORZUN and A. MEISTER, 2000 Cytologically integrated physical restriction fragment length polymorphism maps for the barley genome based on translocation breakpoints. *Genetics* **154**: 397–412.
- KYNAST, R. G., O. RIERA-LIZARAZU, M. I. VALES, R. J. OKAGAKI, S. B. MAQUIEIRA *et al.*, 2001 A complete set of maize individual chromosome additions to the oat genome. *Plant Physiol.* **125**: 1216–1227.
- LAMB, J. C., and J. A. BIRCHLER, 2006 Retroelement genome painting: cytological visualization of retroelement expansions in the genera *Zea* and *Tripsacum*. *Genetics* **173**: 1007–1021.
- LAMB, J. C., T. DANILOVA, M. J. BAUER, J. M. MEYER, J. J. HOLLAND *et al.*, 2007 Single-gene detection and karyotyping using small-target fluorescence *in situ* hybridization on maize somatic chromosomes. *Genetics* **175**: 1047–1058.
- LAWRENCE, C. J., T. E. SEIGFRIED, H. W. BASS and L. K. ANDERSON, 2006 Predicting chromosomal locations of genetically mapped loci in maize using the Morgan2McClintock Translator. *Genetics* **172**: 2007–2009.
- LEE, H. R., W. ZHANG, T. LANGDON, W. JIN, H. YAN *et al.*, 2005 Chromatin immunoprecipitation cloning reveals rapid evolutionary patterns of centromeric DNA in *Oryza* species. *Proc. Natl. Acad. Sci. USA* **102**: 11793–11798.
- LEE, M., N. SHAROPOVA, W. D. BEAVIS, D. GRANT, M. KATT *et al.*, 2002 Expanding the genetic map of maize with the intermated B73 x Mo17 (IBM) population. *Plant Mol. Biol.* **48**: 453–461.
- LIN, Y.-R., L. ZHU, S. REN, J. YANG, K. F. SCHERTZ *et al.*, 1999 A *Sorghum propinquum* BAC library, suitable for cloning genes associated with loss-of-function mutations during crop domestication. *Mol. Breed.* **5**: 511–520.
- LIU, K., M. GOODMAN, S. MUSE, J. S. SMITH, E. BUCKLER *et al.*, 2003 Genetic structure and diversity among maize inbred lines as inferred from DNA microsatellites. *Genetics* **165**: 2117–2128.
- LIU, R., C. VITTE, J. MA, A. A. MAHAMA, T. DHLIWAYO *et al.*, 2007 A GeneTrek analysis of the maize genome. *Proc. Natl. Acad. Sci. USA* **104**: 11844–11849.

- LUCE, A. C., A. SHARMA, O. S. MOLLERE, T. K. WOLFRUBER, K. NAGAKI *et al.*, 2006 Precise centromere mapping using a combination of repeat junction markers and chromatin immunoprecipitation-polymerase chain reaction. *Genetics* **174**: 1057–1061.
- LYSAK, M. A., P. F. FRANSZ, H. B. ALI and I. SCHUBERT, 2001 Chromosome painting in *Arabidopsis thaliana*. *Plant J.* **28**: 689–697.
- MA, J., L. ZHANG, B. B. SUH, B. J. RANEY, R. C. BURHANS *et al.*, 2006 Reconstructing contiguous regions of an ancestral genome. *Genome Res.* **16**: 1557–1565.
- MARTIENSSSEN, R. A., P. D. RABINOWICZ, A. O'SHAUGHNESSY and W. R. MCCOMBIE, 2004 Sequencing the maize genome. *Curr. Opin. Plant Biol.* **7**: 102–107.
- MCCLINTOCK, B., 1941 The stability of broken ends of chromosomes in *Zea mays*. *Genetics* **26**: 234–282.
- MCCLINTOCK, B., 1978 Mechanisms that rapidly reorganize the genome. *Stadler Genet. Symp.* **10**: 25–48.
- MESSING, J., and H. K. DOONER, 2006 Organization and variability of the maize genome. *Curr. Opin. Plant Biol.* **9**: 157–163.
- MESSING, J., A. K. BHARTI, W. M. KARLOWSKI, H. GUNDLACH, H. R. KIM *et al.*, 2004 Sequence composition and genome organization of maize. *Proc. Natl. Acad. Sci. USA* **101**: 14349–14354.
- MEYERS, B. C., S. V. TINGEY and M. MORGANTE, 2001 Abundance, distribution, and transcriptional activity of repetitive elements in the maize genome. *Genome Res.* **11**: 1660–1676.
- NAGAKI, K., Z. CHENG, S. OUYANG, P. B. TALBERT, M. KIM *et al.*, 2004 Sequencing of a rice centromere uncovers active genes. *Nat. Genet.* **36**: 138–145.
- NELSON, W. M., A. K. BHARTI, E. BUTLER, F. WEI, G. FUKS *et al.*, 2005 Whole-genome validation of high-information-content fingerprinting. *Plant Physiol.* **139**: 27–38.
- OHMIDO, N., Y. AKIYAMA and K. FUKUI, 1998 Physical mapping of unique nucleotide sequences on identified rice chromosomes. *Plant Mol. Biol.* **38**: 1043–1052.
- PAMPANWAR, V., F. ENGLER, J. HATFIELD, S. BLUNDY, G. GUPTA *et al.*, 2005 FPC Web tools for rice, maize, and distribution. *Plant Physiol.* **138**: 116–126.
- PATERSON, A. H., M. FREELING and T. SASAKI, 2005 Grains of knowledge: genomics of model cereals. *Genome Res.* **15**: 1643–1650.
- PEDERSEN, C., and I. LINDE-LAURSEN, 1994 Chromosomal locations of four minor rDNA loci and a marker microsatellite sequence in barley. *Chromosome Res.* **2**: 65–71.
- PETERSON, D. G., N. L. LAPITAN and S. M. STACK, 1999 Localization of single- and low-copy sequences on tomato synaptonemal complex spreads using fluorescence *in situ* hybridization (FISH). *Genetics* **152**: 427–439.
- RHOADES, M. M., 1950 Meiosis in maize. *J. Hered.* **61**: 59–67.
- RHOADES, M. M., and B. MCCLINTOCK, 1935 The cytogenetics of maize. *Bot. Rev.* **1**: 292–325.
- ROCCHI, M., N. ARCHIDIACONO and R. STANYON, 2006 Ancestral genomes reconstruction: an integrated, multi-disciplinary approach is needed. *Genome Res.* **16**: 1441–1444.
- SADDER, M., and G. WEBER, 2001 Karyotype of maize (*Zea mays* L.) mitotic metaphase chromosomes as revealed by fluorescence *in situ* hybridization (FISH) with cytogenetic DNA markers. *Plant Mol. Biol.* **19**: 117–123.
- SADDER, M. T., and G. WEBER, 2002 Comparison between genetic and physical maps in *Zea mays* L. of molecular markers linked to resistance against *Diatraea* spp. *Theor. Appl. Genet.* **104**: 908–915.
- SADDER, M. T., N. PONELIES, U. BORN and G. WEBER, 2000 Physical localization of single-copy sequences on pachytene chromosomes in maize (*Zea mays* L.) by chromosome *in situ* suppression hybridization. *Genome* **43**: 1081–1083.
- SENIOR, M., and M. HEUN, 1993 Mapping maize microsatellites and polymerase chain reaction confirmation of the targeted repeats using a CT primer. *Genome* **36**: 884–889.
- SHAROPOVA, N., M. D. MCMULLEN, L. SCHULTZ, S. SCHROEDER, H. SANCHEZ-VILLEDA *et al.*, 2002 Development and mapping of SSR markers for maize. *Plant Mol. Biol.* **48**: 463–481.
- SHEN, D.-L., Z.-F. WANG and M. WU, 1987 Gene mapping on maize pachytene chromosomes by *in situ* hybridization. *Chromosoma* **95**: 311–314.
- SHERIDAN, W. F., and D. L. AUGER, 2006 Construction and uses of new compound B-A-A maize chromosome translocations. *Genetics* **174**: 1755–1765.
- SHERMAN, J. D., and S. M. STACK, 1995 Two-dimensional spreads of synaptonemal complexes from solanaceous plants. VI. High-resolution recombination nodule map for tomato (*Lycopersicon esculentum*). *Genetics* **141**: 683–708.
- SONG, B. K., K. NADARAJAH, M. N. ROMANOV and W. RATNAM, 2005 Cross-species bacterial artificial chromosome (BAC) library screening via overgo-based hybridization and BAC-contig mapping of a yield enhancement quantitative trait locus (QTL) yld1.1 in the Malaysian wild rice *Oryza rufipogon*. *Cell Mol. Biol. Lett.* **10**: 425–437.
- SONG, J., F. DONG and J. JIANG, 2000 Construction of a bacterial artificial chromosome (BAC) library for potato molecular cytogenetics research. *Genome* **43**: 199–204.
- SWIGONOVA, Z., J. LAI, J. MA, W. RAMAKRISHNA, V. LLACA *et al.*, 2004 Close split of sorghum and maize genome progenitors. *Genome Res.* **14**: 1916–1923.
- TARAMINO, G., and S. TINGEY, 1996 Simple sequence repeats for germplasm analysis and mapping in maize. *Genome* **39**: 277–287.
- TRASK, B. J., 2002 Human cytogenetics: 46 chromosomes, 46 years and counting. *Nat. Rev. Genet.* **3**: 769–778.
- WALLING, J. G., R. SHOEMAKER, N. YOUNG, J. MUDGE and S. JACKSON, 2006 Chromosome-level homeology in paleopolyploid soybean (*Glycine max*) revealed through integration of genetic and chromosome maps. *Genetics* **172**: 1893–1900.
- WANG, C. J., L. HARPER and W. Z. CANDE, 2006 High-resolution single-copy gene fluorescence *in situ* hybridization and its use in the construction of a cytogenetic map of maize chromosome 9. *Plant Cell* **18**: 529–544.
- WHITE, S., and J. DOEBLEY, 1998 Of genes and genomes and the origin of maize. *Trends Genet.* **14**: 327–332.
- WOO, S. S., J. JIANG, B. S. GILL, A. H. PATERSON and R. A. WING, 1994 Construction and characterization of a bacterial artificial chromosome library of *Sorghum bicolor*. *Nucleic Acids Res.* **22**: 4922–4931.
- YIM, Y. S., P. MOAK, H. SANCHEZ-VILLEDA, T. A. MUSKET, P. CLOSE *et al.*, 2007 A BAC pooling strategy combined with PCR-based screenings in a large, highly repetitive genome enables integration of the maize genetic and physical maps. *BMC Genomics* **8**: 47.
- YUAN, Y., P. J. SANMIGUEL and J. L. BENNETZEN, 2003 High-Cot sequence analysis of the maize genome. *Plant J.* **34**: 249–255.
- ZHONG, X. B., H. J. DE JONG and P. ZABEL, 1996a Preparation of tomato meiotic pachytene and mitotic metaphase chromosomes suitable for fluorescence *in situ* hybridization (FISH). *Chromosome Res.* **4**: 24–28.
- ZHONG, X. B., P. F. FRANSZ, J. WENNEKES-VAN EDEN, P. ZABEL, A. VAN KAMMEN *et al.*, 1996b High-resolution mapping on pachytene chromosomes and extended DNA fibres by fluorescence *in situ* hybridization. *Plant Mol. Biol. Rep.* **14**: 232–242.
- ZHONG, X. B., J. BODEAU, P. F. FRANSZ, V. M. WILLIAMSON, A. VAN KAMMEN *et al.*, 1999 FISH to meiotic pachytene chromosomes of tomato locates the root-knot nematode resistance gene *Mi-1* and the acid phosphatase gene *Aps-1* near the junction of euchromatin and pericentromeric heterochromatin of chromosome arms 6S and 6L, respectively. *Theor. Appl. Genet.* **98**: 365–370.
- ZWICK, M. S., M. N. ISLAM-FARIDI, D. G. J. CZESCHIN, R. A. WING, G. E. HART *et al.*, 1998 Physical mapping of the liguleless linkage group in *Sorghum bicolor* using rice RFLP-selected sorghum BACs. *Genetics* **148**: 1983–1992.